

PRODUCTION OF CHARGED AND NEUTRAL HIGGS
BOSONS WITH CHARGINOS AND NEUTRALINOS
THROUGH DIFFERENT PROPAGATORS IN THE *MSSM*

BY

HATIM HEGAB ALI HEGAB

A Dissertation submitted to the Faculty of Science, Cairo University
in partial fulfillment of the requirements
for the Degree

Master of Science, Physics.

Major Subject: Physics

Minor Subject: Theoretical High Energy Physics

Cairo University
Giza
Egypt

March, 2006

Copyright 2006 by Hatim Hegab
All rights reserved

“Production of Charged and Neutral Higgs Bosons with Charginos and Neutralinos through different Propagators in the *MSSM*,” a dissertation prepared by Hatim Hegab Ali Hegab in partial fulfillment of the requirements for the degree, Master of Science, Physics. , has been approved and accepted by the following:

Prof. M. M. El-Khishen
Prof. of Theoretical Physics,
Faculty of Science,
Cairo University.

Dr. Tareq Abdul Aziem
Lecturer,
Faculty of Science,
Cairo University.

Date

Committee in charge:

Prof. Dr. M. M. El-Khishen, Chair

Prof. of Theoretical Physics,

Faculty of Science,

Cairo University.

Dr. Tareq Abdul Aziem

lecturer,

Faculty of Science,

Cairo University.

DEDICATION

This work is dedicated to my first teacher of mathematics, **My Mother**.

And to my father, the late **Prof. Hegab Ali Hegab, ex. Prof. of Microbiology at The Technical Agricultural Institute in Kaseem, Bureidah, Saudi Arabia** whose guidance, instructions and kind heart helped to shape and build my personality.

And finally, to my wife, and my daughter, **Basma (A “Smile” in the Arabic Language)**, your existence in my life gave me the power to do much more than I could do alone.

ACKNOWLEDGMENTS

The author would like to thank all those people who helped make this work possible, I here want to express my deepest gratitude to the following people for their help,

1. Prof. M. M. El-Khishen, for his kind supervision and his helpful manners.
2. Dr. Tareq Abdul Aziem.
3. Prof. Alexander Pukhov, for his fruitful discussions, cooperative manners, without his help, I wouldn't be able to do this work.

ABSTRACT

PRODUCTION OF CHARGED AND NEUTRAL HIGGS BOSONS WITH CHARGINOS AND NEUTRALINOS THROUGH DIFFERENT PROPAGATORS IN THE *MSSM*

BY

HATIM HEGAB ALI HEGAB

Master of Science, Physics.

Cairo University

Giza

Egypt, 2006

Dr. Tarek Abdul Aziem

In this work, the following reactions have been studied;

1. $e^-e^+ \rightarrow H^- \tilde{\chi}_1^+ \tilde{\chi}_1^o$

2. $e^-e^+ \rightarrow h \tilde{\chi}_1^+ \tilde{\chi}_1^-$

3. $e^-e^+ \rightarrow h \tilde{\chi}_1^o \tilde{\chi}_1^o$

4. $e^-e^+ \rightarrow h H^+ H^-.$

where $\tilde{\chi}_1^\pm$ is the chargino and $\tilde{\chi}_1^0$ is the neutralino. h represent the lightest neutral Higgs boson, and H_3 (sometimes referred to as A) is the pseudoscalar Higgs boson.

The work went to calculate the differential cross section due to each Feynman diagram, then, the *total cross section* for each reaction is calculated according to a carefully used set of parameters. Results are graphed and tabulated.

CONTENTS

I	Theoretical Framework	xii
1	STANDARD MODEL <i>OF</i> ELECTROWEAK INTERACTIONS	1
1.1	Introduction	1
1.2	Gauge Invariance	3
1.3	Renormalization	5
1.4	Quantum Electrodynamics	6
1.5	Yang-Mills Theory	9
1.6	The Higgs Mechanism	14
1.7	Glashow, Weinberg <i>and</i> Salam Theory	20
2	SUPERSYMMETRY	36
2.1	Motivation <i>for</i> Supersymmetry	36
2.2	Rules <i>of</i> Supersymmetry	41
2.3	Essentials <i>of</i> Supersymmetry Algebra	46
2.4	The Algebra of N=1 supersymmetry	53
2.5	The Wess-Zumino Model	59
2.6	Mass <i>and</i> Interaction Terms <i>in the</i> Lagrangian	65
2.7	The Superpotential	67
2.8	Supersymmetric Gauge Theory	70
2.9	Spontaneous Breaking <i>of</i> Supersymmetry	73

2.10	F-type Breaking (O’raifeartaigh Model)	78
2.11	D-type Breaking (The Fayet-Iliopoulos Model)	82
2.12	The Supersymmetric Standard Model	85
3	MINIMAL SUPERSYMMETRIC STANDARD MODEL	88
3.1	Introduction	88
3.2	Extended Higgs Sectors	92
3.2.1	The ρ –Parameter constraint	93
3.2.2	The $W^-Z^0H^-$ Couplings	94
3.3	The Two Higgs Doublet Model	95
3.4	The Higgs Sector <i>of the</i> MSSM	99
3.5	The Supersymmetric Particle Sector <i>of the</i> MSSM	103
3.5.1	The charginos	105
3.5.2	The Neutralinos	108
3.5.3	The Sferminos	112
3.6	Reducing <i>the</i> MSSM Parameter Freedom	113
3.6.1	Minimization <i>of the</i> Higgs potential	115
3.6.2	Diophantine Analysis <i>of the</i> Higgs Boson	118
3.7	Conclusion	120
II	Phenomenological Calculations	122
4	INTRODUCTION	123

5	PRODUCTION OF A CHARGED HIGGS BOSON WITH A CHARGINO AND A NEUTRALINO	128
5.1	Introduction	128
5.2	Feynman Diagrams	129
5.3	Matrix Elements	132
5.4	Cross Sections	134
5.5	Conclusion	164
6	PRODUCTION OF A LIGHT NEUTRAL HIGGS BOSON WITH A CHARGINO AND A NEUTRALINO	170
6.1	Introduction	170
6.2	Feynman Diagrams	171
6.3	Matrix Elements	173
6.4	Cross Sections	175
6.5	Conclusion	190
7	PRODUCTION OF A LIGHT NEUTRAL HIGGS BOSON WITH A PAIR OF NEUTRALINOS	193
7.1	Introduction	193
7.2	Feynman Diagrams	194
7.3	Matrix Elements	197
7.4	Cross Sections	199

7.5	Conclusion	225
8	PRODUCTION OF A LIGHT NEUTRAL HIGGS BOSON WITH A PAIR OF CHARGED HIGGS BOSONS	230
8.1	Introduction	230
8.2	Feynman Diagrams	231
8.3	Matrix Elements	232
8.4	Cross Sections	233
8.5	Conclusion	240
APPENDICES		
A.	Appendix A: Feynman Rules	242
	Fermions	243
	Vector Bosons	243
	Scalar (Higgs) Bosons	243
	fermions	243
	Massive Vector Bosons	243
	Massless Vector Bosons (Photons)	244
	Scalar (Higgs) Bosons	244
B.	Appendix B: Numerical Values of Constants	245

Part I

Theoretical Framework

Introduction

Progress in theoretical physics, as in all sciences, has almost always been based on an interplay of two quite different approaches of nature.

One starts by collecting and ordering observational, or experimental, data “ Tycho”, then describes these data by a small number of empirical laws “ Kepler”, and finally *explains* these laws by a **theory**, based on a few principles “ **Newton**”. Theoretical predictions, or the outcome, of further, more refined, observations and experiments can then be made (“ discovery of Neptune”).

the other approach starts from an idea, formulates it in terms of a theory, and proceeds to make predictions which then acts as a test of the theory and of its original idea. The later approach – in its pure form– has been most dramatically and singularly successful in Einstein’s development of the Theory of General Relativity (TGR). **Supersymmetry** (**SUSY**) has started from an idea, and at the moment a huge work is going on to confirm this idea.

Modern particle physics, in seeking a single unified theory of all elementary particles and their fundamental interactions, appear to be reaching the limits of this process and finds itself forced, in part and often reluctantly, to revert to guidelines to the medieval principles of symmetry and beauty.

Supersymmetric theories are highly symmetric and very beautiful.

They are remarkable in that they unify fermions (matter) with bosons (force carriers), either in flat space (SUSY) or in curved space, supergravity, (SUGRA). **SuperGravity**,

(**SUGRA**) naturally unifies gravity with other interactions. None of the present model theories is in any sense complete; the hurdles on the way to experimental predictions– and thus to acceptance or rejection– have not yet been cleared. What naive immediate predictions can be made seem to be in disagreement with nature. Yet, this particular field of research appears to promise solutions of so many outstanding problems that it has excited enthusiasm in large parts of the physics community (and equally large doubts in others). In a truly philosophical spirits has been even said about the theory **that it is so beautiful so it is hard to be incorrect.**

In high energy physics, or as it is sometimes called- elementary particle physics, the hope is that we will eventually achieve a unified scheme which combines all particles and all interactions into one consisting theory. We wish to make further progress on the way which started with Maxwell’s unification of the magnetism and electrostatics, and which has more recently led to unified gauge theories (UGT) of electromagnetic and the weak, and perhaps also of the strong interactions.

Supersymmetry is, by definition, a symmetry between fermions and bosons. A supersymmetric field theoretical model consists of a set of quantum fields and of a Lagrangian for them which exhibit such a symmetry. The Lagrangian determines, through the Action Principle, the equations of motion and hence the dynamic behavior of the particles. A supersymmetric model which is covariant under general coordinate transformations, or equally, a model which possesses a local (“ gauged”) supersymmetry is called a supergravity model. Supersymmetric theories describe model worlds of particles, created from

the vacuum by the fields, and the interactions between these particles. The supersymmetry manifests itself in the particle spectrum and in the stringent relationship between different interaction processes even if these processes involve particles of different spin and statistics.

Both supersymmetry and supergravity aim at unified description of *fermions* and *bosons*, and hence of matter and interactions. Supergravity is particularly promising at its attempt to *unify gravity with other interactions*. All supersymmetric models succeed to some extent in these aims, but they fail in describing the real world as we live in and experience it and thus are models not theories. We are still to struggle to find some contact between one of the models and the physical world, reality, so that the model could become an underlying theory for nature and its “most fundamental level”.

By “most fundamental level” we mean at present the decomposition of matter into quarks and leptons (fermions) and the understanding of all forces between them as arising out of four types of basic interactions, namely, **gravitational, weak, electromagnetic, and strong**. These are described in terms of exchanged particles (bosons). The framework within which these building blocks make up a physical theory is *Relativistic Quantum Field Theory*. Seen at this level, “unification” ought to include all four interactions. There is, however, a quantitative and a qualitative difference between the gravitational interaction and the others which has had profound consequences on both the structure of the universe and on our understanding of it.

Unification of gravity with other forces is an illusive goal. Since gravity is always

attractive and a long range force, it would make all complex and large physical objects collapse under their own weight, however, it is not stronger enough at short distances to have any remarkable effect with respect to other forces, [interactions].

This difference in strength, necessary to the universe as we see, has in turn set gravity so far apart from the rest of physics that it is difficult to think of any experiment which could actually test predictions of a unified field theory of all interactions, and even less of one that could provide experimental input into the construction of such a theory. The natural domain of Newtonian gravity and of its modern replacement , Einstein’s TGR, is the world of large distances and massive accumulations of matter, that of the other forces is the world of atoms, nuclei, and elementary particles. A very large Order of magnitude separate them.

The *strong, electromagnetic and weak interactions* are fairly understood today. It has been found that their exchange particles arise naturally in a quantum field theory, if that field theory is required to be locally gauge invariant.

The theory for both gravitation and elementary particles’ interactions are well established within their respective domains. On the submicroscopic level, for the masses and distances involved, the deviation introduced by gravity from the flat Minkowskian metric are so minute that elementary particles, or atoms, can safely be treated as if gravitation does not exist. Any “ **true**”, that is, generally covariant, theory should thus be closely approximated by Lorentz–covariant, non-gravitational theories. We must, however demand of the *true* theory to be mathematically consistent and that it predicts

the correct flat limit. Any quantum theory of gravitation so far fails to do so.

The energy at which gravity and quantum effects become of comparable strength can be estimated from only an expression with a dimension of an energy that can be formed from the constants of nature $\hbar, c, G$,

$$E_{Planck} = c^2 \sqrt{\frac{\hbar c}{G}} \simeq 10^{19} GeV$$

It is in the region of this energy, *Planck's energy*, where our present theories for the gravitational and other interactions become incompatible with each other and where we expect a possible unification of the interactions to become manifest. A point particle with Plank mass would have a Schwarzschild radius equal to twice its Compton wavelength. The very remoteness of such an energy region eliminates all hope for a direct experimental proof. Perhaps, if we are lucky enough, some isolated prediction of such a unified theory could be testable on a system that integrates a minute effect over a vast range (proton decay experiments are of this type, where large numbers of available protons can make the very small probabilities for decay available and measurable). We can, however, not expect experimental physics to give us much reliable guidance into the *Plank region*.

The $SU(3) \times SU(2) \times U(1)$ picture of the non-gravitational forces is not yet a *unified* one. The only property which unites them is that they are each described by a gauge theory. The fact that the direct product structure of $SU(2) \times U(1)$ is “skew” (by the weak mixing angle) against the natural distinction between the weak and electromagnetic

force may suggest some underlying unified scheme which can predict that angle. A lot of work has, over many years, gone into finding a larger gauge group which would describe all three interactions at some high energy. If such a grand unification occurred it is known that it must happen at energies of about $10^{15} GeV$, only four orders of magnitude less than E_{Planck} . *Grand Unified Theories* (GUTs) have had some success (such as the prediction of the mixing angle) and some failures (such as proton life time experiments). In any case, even a GUT would at most unify different kinds of interactions (strong and electroweak) with each other and different kinds of matter (quarks and leptons) with each other. the unification of matter with interactions is not one of the aims of GUTs.

What is it then that points in the direction of supersymmetric theories for a *solution to the unification problem*? Already the most obvious difference between gravity and, say, electrodynamics, namely the absence of negative gravitational charges, can be shown to imply that only a supersymmetric theory can unify them. As long as we do not dramatically deviate from standard quantum field theory, and we hope that, that will not be necessary, a purely attractive force must be carried by a field with even integer spin. The electromagnetic force, on the other hand, which is- of course- not always attractive, is carried by a field of spin one. A number of no-go theorems about which we will have to know more later, forbids any direct symmetry transformations between fields of different integer spin and actually leave supersymmetric theories as the only fields theoretical models which achieve unification of all forces of nature. Supersymmetry transformations do not directly relate fields of different integer spin, rather they relate a

graviton (a quantum of gravity with spin 2) to a photon (spin 1) via a spin $\frac{3}{2}$ intermediary, the *gravitino*.

A partial unification of matter (fermions) with interactions (bosons) thus arises naturally out of the attempt to unite gravity with the other interactions.

The no-go theorems imply that *supersymmetry* and *supergravity* are only possibilities for unification within the framework of quantum field theory. Failure of these theories to ultimately give results which are compatible with the real world would force us to give up either unification or quantum field theory.

A part from the no-go theorems there is a further, more technical point that singles out supersymmetric theories; they may resolve the nonrenormalizability problem of quantized gravity. In perturbation quantum field theory fermions and bosons tend to contribute with opposite signs to higher order corrections. Supersymmetry almost always achieve a fine-tuning between these contributions which makes some a priori present and *divergent terms vanish*. For a long time now there has been great optimism that of the supergravity models may be entirely free of infinities and thus be a consistent model of quantized gravitation with no need for renormalization. This hope is slowly fading now, but only to the extent that no obvious proof can be found for the conjectured finiteness. What remains is a remarkably friendly behavior of all supersymmetric theories when it comes to quantum divergences, and the conviction of many theorists that it would be surprising if nature did not make use of it. We will have to say later more about *cancellations of divergences* and the enormous interest which

they have aroused, not only for gravity but also for the *hierarchy problem* of GUTs, the thirteen orders of magnitude between the GUT mass of $10^{15} GeV/c^2$ and the W -boson mass. Normally, a gap of this size is not stable in perturbation theory because of the considerable admixture of the large mass to the small one through vacuum polarization. the gap can only be maintained by repeated fine-tuning up to high orders in the perturbation expansion. In supersymmetric versions of GUTs new particles are exchanged and pair-created, and these new processes cancel some of the effects of the old ones. Mass mixing and consequently fine-tuning can usually be avoided, and the hierarchy, once established, is stabilized.

Outline of Part One:

This part consists of four chapters of which the fourth chapter is the main one.

In the **first chapter**, Introduction, an introduction to the subjected presented in this part is given (this one).

In the **second chapter**, The *Standard Model of Electroweak Interactions* is presented with a brief background examples of gauge theories and Higgs mechanism.

In the **third chapter**, The *Supersymmetry concepts, Supersymmetry Algebra, and Supersymmetric Models* are introduced.

And in the **fourth chapter**, The *Minimal Supersymmetric Standard Model* was studied, including *the extended Higgs model* and *the particle spectrum of the model*.

CHAPTER 1

STANDARD MODEL *OF* ELECTROWEAK INTERACTIONS

1.1 Introduction

THE invention of unified renormalization theories of electroweak interactions is actually one of the outstanding successes of elementary particle physics. the first of this theories was the theory of Glashow, Weinberg and Salaam (**GWS**) and is known as the standard electroweak theory.

In 1961 Glashow constructed a model for the weak and electromagnetic interactions of leptons which was based on the assumption that, together with the photon, there exist also a charged W and a neutral Z intermediate bosons. The mass of the W and Z bosons were introduced “by hand”, *ad hoc*. As a result, the model was unrenormalizable. In 1967-1968 Weinberg and Salam constructed the $SU(2) \times U(1)$ model of electroweak interactions of leptons including a spontaneous breakdown of the gauge symmetry. In 1971-1972 it was proved by t’Hooft and others that the model of this type were renormalizable. The model was generalized to quarks using the mechanism proposed by Glashow, Iliopoulos, and Maiani.

The GS theory is based on the assumption of the existence of charged and neu-

tral intermediate vector bosons and it is constructed so that, for massless fundamental fermions (leptons and quarks), a local $SU(2) \times SU(1)$ gauge invariance takes place. then the interaction (again locally gauge invariance) of Higgs scalar fields with both gauge vector bosons and fermions, is introduced. As a consequence of the spontaneous breakdown of the underlying symmetry, leptons, quarks, and intermediate bosons all acquire masses.

The only free parameter, which enters in the definition of the neutral current in the GWS theory, is $\sin^2 \theta_W$ (where θ_W is the Weinberg angel)

Neutral currents were discovered at CERN in 1973 in an experiment using the large bubble chamber “Gargamelle”. In this experiment the process $\bar{\nu}_\mu + e \rightarrow \bar{\nu}_\mu + e$ was observed. After the pioneering work of the “Gargamelle” collaboration, a large number of experiments were done investigating various neutral current induced processes. After this work it became possible to perform a complete phenomenological analysis of all the neutral current data. As a result one could uniquely determine all the coefficients appearing in the most general phenomenological V, A expressions written for hadron and lepton neutral currents. It was shown that this unique solution is in agreement with the GWS theory.

In 1980-1981, in experiments on the e^-e^+ colliding beams, information has been obtained on the contribution of neutral currents to the cross sections of the process $e^+ + e^- \rightarrow l^+ + l^- (l = e, \mu, \tau)$. These data also agreed with the *standard electroweak model*.

The *GBS* theory predicts the values of the charged (W) and neutral (Z) intermediate boson masses, namely, $m_W \sim 80\text{GeV}$ and $m_Z \sim 91\text{GeV}$.

the discovery in 1983 of the W and Z bosons at the *CERN* $p\bar{p}$ collider, with exactly the predicted masses, was a dramatic confirmation of the *GBS* theory.

In the *GBS* theory $\sin^2 \theta_W$ is a free parameter. It is related to the value of W and Z masses as $\sin^2 \theta_W = 1 - (\frac{m_W^2}{m_Z^2}) \sim 0.23$

1.2 Gauge Invariance

The concept of gauge invariance [1] grew out of the observation that of a “charge” (e.g. electric charge, total energy, isospin, etc.) is conserved in a dynamical system, then the Lagrangian for the system is invariant under “global gauge transformation” of the fields. For example, the electric charge is related to invariance under phase transformations $\psi \rightarrow e^{iq\theta}\psi$ for all fields ψ which describe particles of charge q . Similarly, the energy is related to time translations $\psi(t, x) \rightarrow \psi(t + \Delta t, x)$. The converse is also true (Nether’s theorem); if the Lagrangian is invariant under some infinitesimal transformation $\psi \rightarrow \psi + \delta\psi$, then there is a conserved current and a conserved charge associated with this gauge invariance, (“gauge” is an unfortunate naming, originating in an attempt by H. Wyle in 1918 to relate the electric charge to re-scaling transformation $\psi \rightarrow e^\lambda\psi$). We call the transformation global if their parameters do not depend on the space-time coordinates, i.e. if $\theta = \text{cons.}$ This relationship between conserved quantum numbers and global symmetries of the Lagrangian led, in the 1960’s, to a search for

globally-invariant field theories capable of describing and classifying all elementary particles. The “ **8-fold way**” very much in this vein and it was in this context that quarks were first postulated as building blocks of strongly interacting particles.

The requirement of the *local gauge invariance* (also known as “ gauge invariance of the 2nd kind”) goes beyond that which can be inferred from charge conservation. We now demand invariance of the Lagrangian under transformations with a space-time dependent parameter $\theta = \theta(x)$. This interaction, which results in the exchange of the field quanta, will generate forces between the particles. “ Gauging” the phase transformation associated with electric charge (i.e. making them *x – dependent*) forces us to introduce the electromagnetic four-vector potential and, as its quanta, the photons. The result is quantum electrodynamics. Requiring other gauge invariances requires additional gauge potentials which give rise to more exchange particles and the other interactions. These exchanged particles are the discovered $W^\pm$ and Z^0 for the weak force and the gluons for the strong interactions. The latter have only been indirectly seen in their effects on the distribution of the debris in high energy particle collisions (*jets*). To sum up, “ gauging” an invariance of the Lagrangian will always give rise to interaction and to forces.

Nowadays, the name gauge theory is used exclusively for theories with local gauge invariance.

The gauge transformation under which the Lagrangian is invariant forms a group as it fulfills the axioms of a group (in the mathematical sense):

- 1- Two subsequent invariances will again be an invariance,

- 2- “No transformation” is the identity element,
 - 3- There is exactly one inverse transformation for each invariance transformation,
- and
- 4- Three transformations are associative.

Using the standard terminology of groups, the respective gauge groups are $SU(3)$ for the strong interactions and $SU(2) \times U(1)$ for the electroweak interactions. The $SU(3)$ transformations act on triplets of quarks whose properties are very similar. They are said to differ only in “**color**”, hence the name quantum-chromodynamics (*QCD*) for the $SU(3)$ gauge theory of the strong interactions. The success of gauge theories in describing a variety of elementary particles phenomena eclipsed the rule played by global invariance, and nowadays such global symmetries are thought of as more or less accidental -if indeed they are present. In this context it is already important to mention that local (gauged) supersymmetry will imply supergravity.

1.3 Renormalization

Renormalization is required in all quantum field theories in order to make sense of divergent integrals which appear in the perturbation expansions for physical processes. Such expansions are unfortunately the only calculations tools currently available for solving the equations of motion of the theory; they are usually conceptualized in terms of vacuum polarizations and virtual particle interactions and are illustrated by Feynman Diagrams. In renormalizable theories, the divergences which appear can be tested by

redefining, in each order of the perturbation expansion, a finite number of theoretical parameters in such a way that the results of “test experiments” are reproduced. Other processes can be calculated uniquely to the same order. In the lowest order, the parameters which must be so renormalized typically present vacuum energies, masses, coupling constants and factors which are multiplied to the wave functions. Correspondingly, one speaks of “vacuum, mass, coupling constant, and wave function renormalization”. One of the strongest motivations for gauge theories is their renormalizability.

A theory is called non-renormalizable if infinitely many parameters must be redefined. Such a “theory” can make no predictions and therefore not a theory in the sense of exact science. In general, coupling constants with negative mass dimensions (for $\hbar = c = 1$) lead to non-renormalizable theories. No matter how we attempt to quantize gravity, we end up with a field theory whose coupling constant, **Newton’s universal gravitational constant** G , has dimensions $\frac{1}{mass^2}$ in these units and quantum gravity is therefore non-renormalizable.

1.4 Quantum Electrodynamics

Quantum Electrodynamics (QED) is the gauge invariance theory which describes all relevant experimental data. As an example, consider the electron field $\psi(x)$. The free Lagrangian of this field has the standard form,[2].

$$\mathcal{L} = -\bar{\psi}(\gamma_\alpha \partial_\alpha + m)\psi \tag{1.1}$$

where m is the mass of the electron, $\partial_\alpha \equiv \frac{\partial}{\partial x_\alpha}$. The Lagrangian (1.1) is invariant with respect to the global gauge transformation

$$\psi(x) \rightarrow \psi'(x) = e^{i\lambda} \psi(x), \quad (1.2)$$

where λ is an arbitrary real constant. It is obvious that the Lagrangian (1.1) is not invariant with respect to the local gauge transformation

$$\psi(x) \rightarrow \psi'(x) = U(x) \psi(x) \quad (1.3)$$

where

$$U(x) = \exp\{i\lambda(x)\}$$

and where $\lambda(x)$ is an arbitrary real function of x . The derivative $\partial_\alpha \psi(x)$ is indeed not transformed under (1.3) as the field $\psi(x)$ itself. Really, we have

$$\partial_\alpha \psi'(x) = U(x) (\partial_\alpha + i\partial_\alpha \lambda(x)) \psi(x)$$

As is well known, the local gauge invariance (1.3) can be maintained provided that the interaction of the field ψ with electromagnetic field A_α is introduced. Consider the quantity $(\partial_\alpha - ieA_\alpha)\psi$ (e is the electron charge), we will have

$$(\partial_\alpha - ieA_\alpha(x))\psi(x) = U^{-1}(x)(\partial_\alpha - ieA'_\alpha(x))\psi'(x)(\partial_\alpha - ieA_\alpha)\psi \quad (1.4)$$

where

$$A'(x) = A_\alpha(x) + \frac{1}{e}\partial_\alpha\lambda(x) \quad (1.5)$$

From (1.4) it is obvious that the Lagrangian, which follows from (1.1) by the substitution

$$\partial_\alpha\psi \rightarrow (\partial_\alpha - ieA_\alpha)\psi \quad (1.6)$$

is now invariant with respect to the gauge transformation (1.3) and (1.5).

To construct the complete Lagrangian of the system under consideration, we have to add also the gauge invariant Lagrangian of the electromagnetic field. the tensor of the electromagnetic field is given as

$$F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha \quad (1.7)$$

Clearly, $F'_{\alpha\beta} = F_{\alpha\beta}$, consequently, the gauge invariant Lagrangian of the fields of electrons and photons takes the form

$$\mathfrak{L} = -\bar{\psi}[\gamma_\alpha(\partial_\alpha - ieA_\alpha) + m]\psi - \frac{1}{4}F_{\alpha\beta}F_{\alpha\beta} \quad (1.8)$$

The substitution of the derivative $\partial_\alpha\psi$ by the covariant derivative $(\partial_\alpha - ieA_\alpha)\psi$ in the free Lagrangian of the field ψ leads to the following interaction Lagrangian for electrons and photons;

$$\mathfrak{L}_i = iej_\alpha A_\alpha \quad (1.9)$$

where $j_\alpha = \bar{\psi}\gamma_\alpha\psi$ is the electromagnetic current. Thus the substitution (1.6) fixes uniquely the form of the interaction Lagrangian. Such an interaction is called minimal electromagnetic interaction. Let us note however that the principle of gauge invariance alone does not fix the interaction Lagrangian uniquely. For example, the addition of the Pauli term $\mu\bar{\psi}\sigma_{\alpha\beta}\psi F_{\alpha\beta}$ to the Lagrangian (1.8) does not spoil the gauge invariance of the theory, (μ is the anomalous magnetic moment).

All available experimental data confirm that the Lagrangian (1.9) is the true Lagrangian which governs the interactions of electrons and photons. It is also well known that electrodynamics, with the minimal interaction (1.9), is a renormalizable theory.

1.5 Yang-Mills Theory

The modern theory of weak interactions is constructed in analogy with quantum electrodynamics. We know from the experiment that the Hamiltonian of weak interactions contains charged currents. Therefore, to construct a theory of weak interactions we have to start with a gauge theory containing fields of charged vector particles. Such a theory does exist. It is the **Yang-Mills** theory which we will now briefly present.

consider the doublet

$$\psi = \begin{pmatrix} \psi^{(1)} \\ \psi^{(-1)} \end{pmatrix}$$

of the group $SU(2)$ ($\psi^{(1)}, \psi^{(-1)}$ are spinor fields). The Lagrangian of the field ψ is written as

$$\mathfrak{L}_0 = -\bar{\psi}(\gamma_\alpha \partial_\alpha + m)\psi \quad (1.10)$$

where m is the common mass of particles, which corresponds to the fields $\psi^{(1)}, \psi^{(-1)}$. Obviously, The Lagrangian (1.10) is left invariant with respect to the global $SU(2)$ transformation

$$\psi(x) \rightarrow \psi'(x) = \exp\{i\frac{1}{2}\tau\lambda\}\psi(x) \quad (1.11)$$

Here τ_i are Pauli matrices and λ_i are real constants.

We are now interested in the conditions under which the Lagrangian of the system is invariant with respect to the local $SU(2)$ transformations

$$\psi(x) \rightarrow \psi'(x) = U(x)\psi(x) \quad (1.12)$$

where

$$U(x) = \exp\{i\frac{1}{2}\tau\lambda(x)\}$$

and where $\lambda(x)$ are arbitrary real functions of x . It is sufficient to consider only the infinitesimal transformations (1.12). The parameters λ_i will be taken as infinitesimal and in all expansions in powers of λ_i we shall keep only the linear terms. Thus we have

$$U(x) \cong 1 + i\frac{1}{2}\tau\lambda(x) \quad (1.13)$$

Next, we get

$$\partial_\alpha \psi(x) = U^{-1}(x) \left(\partial_\alpha - i \frac{1}{2} \tau \lambda(x) \right) \psi'(x) \quad (1.14)$$

It is clear from equation (1.14) that the Lagrangian (1.10) is not invariant under the transformation of (1.12). To construct a gauge invariant theory in analogy with electrodynamics, we thus introduce, besides the field ψ , the vector field A_α . consider the quantity

$$\left(\partial_\alpha - ig \frac{1}{2} \tau A_\alpha \right) \psi(x) \quad (1.15)$$

where g is a dimensionless constant. Using equation (1.13) and the commutation relations $\left[\frac{1}{2} \tau_i, \frac{1}{2} \tau_j \right] = i \epsilon_{ijk} \frac{1}{2} \tau_k$, we find

$$\begin{aligned} \left(\partial_\alpha - ig \frac{1}{2} \tau A_\alpha \right) \psi(x) &= U^{-1}(x) U(x) \left(\partial_\alpha - ig \frac{1}{2} \tau A_\alpha \right) U^{-1}(x) \psi'(x) \\ &= U^{-1}(x) \left(\partial_\alpha - ig \frac{1}{2} \tau A'_\alpha(x) \right) \psi'(x), \end{aligned} \quad (1.16)$$

with

$$A'_\alpha(x) = A_\alpha(x) + \frac{1}{g} \partial_\alpha \lambda(x) - \lambda(x) \times A_\alpha(x) \quad (1.17)$$

The field A_α is called a *Yang – Mills field*. It is seen from equation (1.17), that under the global $SU(2)$ transformations the fields A_α transforms as a triplet.

Thus, as it is seen from the expression (1.16), the covariant derivative $\partial_\alpha - ig\frac{1}{2}\tau A_\alpha$ applied to the field ψ , transforms under the gauge transformations (1.12) and (1.17) as the field ψ itself (a primed quantity is obtained from unprimed one through its multiplication by the matrix U). This means that the substitution for the derivative $\partial_\alpha\psi$ in the free Lagrangian by the covariant derivative (1.15) leads to a Lagrangian, which is invariant with respect to the gauge transformations (1.12) and (1.17).

To construct the gauge invariant Lagrangian of the field A , consider the quantity

$$F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha + gA_\alpha \times A_\beta \quad (1.18)$$

With the help of equation (1.17) it is easy to check that

$$F'_{\alpha\beta} = F_{\alpha\beta} - \lambda \times F_{\alpha\beta} \quad (1.19)$$

It is immediately seen that the quantity $F_{\alpha\beta}F_{\alpha\beta}$ is a group scalar. In analogy with electrodynamics we take the Lagrangian of the field A_α in the form

$$\mathfrak{L}'_0 = -\frac{1}{4}F_{\alpha\beta}F_{\alpha\beta} \quad (1.20)$$

Thus, if the interaction of the fields ψ and A_α is introduced through the “ minimal” substitution $\partial_\alpha\psi \rightarrow (\partial_\alpha - ig\frac{1}{2}\tau A_\alpha)\psi$, the total Lagrangian of the system under consideration has the form

$$\mathfrak{L} = -\bar{\psi} \left[\gamma_\alpha \left(\partial_\alpha - ig\frac{1}{2}\tau A_\alpha \right) + m \right] \psi - \frac{1}{4}F_{\alpha\beta}F_{\alpha\beta} \quad (1.21)$$

Consequently, the interaction Lagrangian of the fields ψ and A_α is as follows:

$$\mathfrak{L}_i = ig\bar{\psi}\gamma_\alpha\frac{1}{2}g\tau\psi A_\alpha \quad (1.22)$$

The constant g introduced before becomes the interaction constant. Therefore the “minimal” substitution $\partial_\alpha\psi \rightarrow (\partial_\alpha - ig\frac{1}{2}\tau A_\alpha)\psi$ fixes uniquely the interaction Lagrangian of the fields ψ and A_α . We have arrived at the “minimal” interaction Lagrangian for the fields ψ and A_α , which is compatible with gauge invariance. One should notice also that owing to the non-linear term $gA_\alpha \times A_\beta$ which appears in the expression (1.18) written for the field tensor $F_{\alpha\beta}$, the Lagrangian (1.21) contains terms which are responsible for the self-interaction of the field A_α .

Notice that a mass term $-\frac{1}{2}m_\gamma^2 A_\alpha A_\alpha$ for the gauge field can not be added to the Lagrangian of the fields of electrons and photons because its presence would destroy the gauge invariance of the theory. this means that the mass of the photon is equal to zero. In this case of the Yang-Mills theory, the imposed gauge invariance also does not allow a mass term of the $-\frac{1}{2}m_\gamma^2 A_\alpha A_\alpha$. Consequently, the particles of the fields A_α are all massless.

We conclude this section with the following remark. Consider several fields $\psi_i (i = 1, 2, 3, \dots, n)$ interacting with the gauge field A_α . We can write

$$\psi'_i(x) = \exp\{i\lambda_i(x)\}\psi_i(x) \quad (1.23)$$

and

$$(\partial_\alpha - ieA_\alpha(x)) \psi(x) = \exp\{i\lambda_i(x)\} (\partial_\alpha - ieA_\alpha(x)) \psi'_i(x) \quad (1.24)$$

where

$$A'_\alpha(x) = A_\alpha(x) + \frac{1}{e_i} \partial_\alpha \lambda_i(x) \quad (1.25)$$

e_i are constants of interaction between the fields ψ_i and the gauge fields A_α . It is clear from equation (1.25) that the local gauge invariance is guaranteed provided that

$$\lambda_i(x) = e_i \lambda(x) \quad (1.26)$$

($\lambda(x)$ is an arbitrary real function of x). Gauge invariance does not impose any restriction on the coupling constants e_i .

In a non-Abelian *Yang – Mills* theory the situation is completely different. If there are several field multiplets interacting with one *Yang – Mills* gauge field, the coupling constants of all the fields with the gauge field are unique. It follows immediately from the fact that the coupling constants enters into the expression for the field tensor $F_{\alpha\beta}$ (eq. (1.18)) because of the non-Abelian character of the Yang-Mills group.

1.6 The Higgs Mechanism

The Lagrangian mass terms are introduced into the *GWS* theory via the so called *Higgs* mechanism for the spontaneous breakdown of the gauge symmetry. To illustrate

how this mechanism works, we shall consider in this section some classical examples of spontaneous symmetry breakdown in relativistic field theory.

Consider for instance the complex scalar field $\phi(x)$ with the Lagrangian density[3]

$$\mathfrak{L} = -\partial_\alpha \phi^* \partial_\alpha \phi - V(\phi^* \phi) \quad (1.27)$$

where

$$V(\phi^* \phi) = -\mu^2 \phi^* \phi + \lambda (\phi^* \phi)^2 \quad (1.28)$$

and where μ^2 and λ are positive constants. The Hamiltonian density obtained from equation (1.27) reads:

$$\mathfrak{H} = \partial_0 \phi^* \phi + \nabla \phi^* \cdot \nabla \phi + V(\phi^* \phi) \quad (1.29)$$

We now look for the minimum of the energy of the system. Obviously, The Hamiltonian (1.29) is minimal at $\phi = \text{const.}$, a value obtained from the condition

$$\frac{\partial V}{\partial \phi} = \phi^* (-\mu^2 + 2\lambda \phi^* \phi) = 0$$

Then we find that the energy of the field is minimal at

$$|\phi_0|^2 = \frac{\mu^2}{\lambda} = \frac{v^2}{2} \quad (1.30)$$

i.e

$$\phi_0 = \frac{v}{\sqrt{2}} e^{i\alpha} \quad (1.31)$$

where α is an arbitrary real parameter. Thus the minimum of the Hamiltonian (1.29) is infinitely degenerate. The degeneracy is obviously connected with the fact the Lagrangian (1.27) is invariant with respect to the global $U(1)$ transformations

$$\phi(x) \rightarrow \phi'(x) = e^{i\lambda} \phi(x) \quad (1.32)$$

The energy minimum of the system under consideration corresponds to an arbitrary value of α in equation (1.31). Due to the gauge invariance of equation (1.32) it is always possible to take

$$\phi_0 = \frac{v}{\sqrt{2}} \quad (1.33)$$

This is the typical example of the spontaneously broken symmetry; the Lagrangian of the field ψ is invariant with respect to the global $U(1)$ transformations, while the value of the field ϕ is invariant with respect to the global $U(1)$ transformations, while the value of the field ϕ , corresponding to the minimal energy, is just one of many possible choices.

We further introduce two real fields χ_1 and χ_2 as

$$\phi = \frac{v}{\sqrt{2}} + \frac{1}{\sqrt{2}}(\chi_1 + i\chi_2) \quad (1.34)$$

It follows from equation (1.33) that the energy of the system reaches its minimum value when the fields χ_1, χ_2 have vanishing values. Substituting (1.34) into equation (1.27), and omitting the unimportant constant $\frac{\lambda v^4}{4}$, we get the Lagrangian of the system in the following form:

$$\mathcal{L} = -\frac{1}{2}\partial_\alpha\chi_1\partial_\alpha\chi_1 - \frac{1}{2}\partial_\alpha\chi_2\partial_\alpha\chi_2 - \frac{1}{4}\lambda(4v^2\chi_1^2 + 4v\chi_1^3 + \chi_1^4 + 4v\chi_1\chi_2^2 + 4\chi_1^2\chi_2^2 + \chi_2^2) \quad (1.35)$$

It now describes the interactions of two neutral scalar fields. The mass term of the field χ_1 is

$$2\lambda v^2\chi_1^2 = m_{\chi_1}^2\chi_1^2 \quad (1.36)$$

Consequently, in the case of quantized fields, the mass of the field quantum χ_1 equals $m_{\chi_1} = \sqrt{2\lambda\mu} = \sqrt{2}\mu$. There is no term quadratic in the field χ_2 . This means that the particle corresponding to the quantum of the field χ_2 is massless.

We have assumed that the value of the constants λ and μ^2 in the Lagrangian (1.27) are positive. Consequently, the quadratic in the field ϕ appears in equation (1.27) are positive, i.e. “wrong” sign. This leads to the spontaneous breaking of the symmetry. The degeneracy of the ground state is a characteristic of this phenomenon. We however introduced new real fields χ_1 and χ_2 for which the ground state is not degenerate. This leads to the spontaneous breakdown of the original $U(1)$ global symmetry of the Lagrangian. As a result, the quanta of one field are massive, while the mass of the second field is zero.

With spontaneous breakdown of continuous symmetry massless spinless (spin zero) particles always appear. This statement is quite general, and it comprises the content of the goldstone theorem. the corresponding massless particles are not observed. This might imply that the ideas of spontaneous symmetry breakdown are useless in constructing realistic physical theories in elementary particle physics. However, it will be shown in the following, how the spontaneous breakdown of local *gauge* symmetry results in massive gauge quanta due to the disappearance of Goldstone bosons.

Let us assume that the complex field ϕ with the Lagrangian (1.27) interacts minimally with the gauge field A_α . This interaction is introduced by the substitution $\partial_\alpha \phi \rightarrow (\partial_\alpha - igA_\alpha) \phi$ in equation (1.27)

The complete Lagrangian of the system is

$$\mathfrak{L} = (\partial_\alpha + igA_\alpha) \phi^* (\partial_\alpha - igA_\alpha) \phi - V(\phi^* \phi) - \frac{1}{4} F_{\alpha\beta} F_{\alpha\beta} \quad (1.37)$$

where

$$F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha \quad (1.38)$$

the Lagrangian (1.37) is invariant with respect to the local gauge transformations

$$\begin{aligned} \phi(x) &\rightarrow \phi^*(x) = e^{i\lambda(x)} \phi, \\ A_\alpha(x) &\rightarrow A'_\alpha = A_\alpha(x) + \frac{1}{g} \partial_\alpha \lambda(x), \end{aligned} \quad (1.39)$$

where $\lambda(x)$ is an arbitrary function of x .

As in the previous example, the minimum of the energy corresponds to a value of the field ϕ equal to $\left(\frac{v}{\sqrt{2}}\right) e^{i\alpha}$ (where α is an arbitrary parameter, $\left(\frac{v}{\sqrt{2}} = \sqrt{\frac{\mu^2}{2\lambda}}\right)$). Due to the gauge invariance of the Lagrangian (1.37) the “vacuum” value of the field ϕ can always be taken as

$$\phi_0 = \frac{v}{\sqrt{2}} \quad (1.40)$$

We shall write the field ϕ in the form

$$\phi(x) = \frac{1}{\sqrt{2}} (v + \chi(x)) e^{i\frac{\theta(x)}{v}} \quad (1.41)$$

where $\chi(x)$ and $\theta(x)$ are real functions of x defined so that zero values correspond to the minimum of V .

It is clear that due to the local gauge invariance of the theory the function $\theta(x)$ appearing in equation (1.41) has no physical meaning. It can always be eliminated by an appropriate gauge transformation. Thus, we have

$$\phi(x) = \frac{(v + \chi(x))}{\sqrt{2}} \quad (1.42)$$

Substituting (1.42) and equation (1.37) and omitting the unimportant constant, we get the Lagrangian of the system under consideration in the following form

$$\mathfrak{L} = -\frac{1}{2}\partial_\alpha\chi\partial_\alpha\chi - \frac{1}{2}g^2(v + \chi)^2 A_\alpha A_\alpha - \frac{1}{4}\lambda(\chi + 2v)^2 \chi^2 - \frac{1}{4}F_{\alpha\beta}F_{\alpha\beta} \quad (1.43)$$

The Lagrangian (1.43) contains the term of the vector field $A_\alpha \left(-\frac{1}{2}g^2v^2A_\alpha A_\alpha\right)$ and the mass term of the scalar field $\chi \left(-\frac{1}{2}2\lambda v^2\chi^2\right)$. Consequently, the masses of the fields A_α and χ are equal to $m_A = gv$ and $m_\chi = \sqrt{2\lambda v^2} = \sqrt{2}\mu$, respectively.

Before spontaneous symmetry breakdown the Lagrangian of the system contained a complex scalar field (two real functions) and a massless gauge field (two independent real functions). After spontaneous breakdown of the local symmetry we arrived at the Lagrangian of an interacting real massive scalar field (one real function) and a massive vector field (three real functions). The degree of freedom, which would correspond to the massless Goldstone boson (in the absence of the gauge field A_α), has been transformed through the spontaneous breakdown of the local gauge symmetry of the Lagrangian (1.37), into the additional degree of freedom (masses of the vector field).

The mechanism thus discussed is called Higgs mechanism. The scalar particle, corresponding to the quantum of the field χ , is called the Higgs particle.

We have explained the basic principles which are used in constructing models of electroweak interactions. Now we turn to the detailed discussion of the standard $SU(2) \times U(1)$ theory of *Glashow, Weinberg, and Salam*.

1.7 Glashow, Weinberg and Salam Theory

The phenomenological V-A (*current* $\times$ *current*)theory was capable of describing the vast amount of existing experimental data. Consequently, any new theory of weak interactions has to be built up as to reproduce the results of the results of this theory.

The GWS theory[4],[5], [6] is based on the assumption that there exists intermediate vector bosons. To reproduce the results of the V-A theory at low energies it is therefore necessary to assume that at least part of the “ true” weak interaction Lagrangian is of the form[7][8]

$$\mathfrak{L} = \frac{ig}{2\sqrt{2}} j_{\alpha}^{(+)} W_{\alpha} + h.c \quad (1.44)$$

where W_{α} is the field of the vector bosons and $j_{\alpha}^{(+)}$ is the charged weak currents.

The dimensionless coupling constant g is related to the fermion constant by

$$\frac{g^2}{8m_W^2} = \frac{G_F}{\sqrt{2}} \quad (1.45)$$

where m_W is the mass of the charged intermediate boson. The charged current is the sum of lepton and hadron (quark) current. In this study we shall consider the *GWS* theory of leptons¹. Consequently, we will be interested only in the lepton current. It follows from all available data that the charged lepton current is

$$j_{\alpha}^{(+)} = \bar{v}_e \gamma_{\alpha} (1 + \gamma_5) e + \bar{v}_{\mu} \gamma_{\alpha} (1 + \gamma_5) \mu + \bar{v}_{\tau} \gamma_{\alpha} (1 + \gamma_5) \tau \quad (1.46)$$

where e, μ, τ are the field operators of the electron, muon and τ -lepton., respectively; v_e, v_{μ}, v_{τ} are the field operators of the electron-, muon-, and tau-neutrinos, respectively.

At the beginning we shall consider the case of massless fields. In order to get the term (1.44) in the interaction Lagrangian of the leptons and vector bosons we assume

¹The study for the case of quarks can be found in reference [2] and the references therein

that

$$\psi_{lL} = \begin{pmatrix} v'_{lL} \\ l'_L \end{pmatrix}, \quad (l = e, \mu, \tau) \quad (1.47)$$

forms a doublet of the $SU(2)$ group and

$$l'_R, v'_{lR} \quad (1.48)$$

are singlets of this group². Here

$$\begin{aligned} \psi_{lL} &= \frac{1}{2}(1 + \gamma_5) \begin{pmatrix} v'_i \\ l' \end{pmatrix} \\ l'_R &= \frac{1}{2}(1 - \gamma_5)l', \\ v'_{lR} &= \frac{1}{2}(1 - \gamma_5)v'_i \end{aligned} \quad (1.49)$$

are the left-handed (L) and the right-handed (R) components of the corresponding fields.

The free field lepton Lagrangian

$$\mathfrak{L} = \sum_{l=e,\mu,\tau} \left(\bar{\psi}_{lL} \gamma_\alpha \partial_\alpha \psi_{lL} + \bar{l}'_R \gamma_\alpha \partial_\alpha l'_R + \bar{v}'_{lR} \gamma_\alpha \partial_\alpha v'_{lR} \right) \quad (1.50)$$

is clearly invariant with respect to the global $SU(2)$ group. We demand now for massless fields the local *Yang – Mills* invariance with respect to

²the primes put on lepton fields indicates that these fields do not necessarily correspond to lepton fields with well defined masses which will be generated later through spontaneous breakdown of the underlying symmetry.

$$\psi_{lL}(x) \rightarrow \psi'_{lL}(x) = \exp\{i\frac{1}{2}\tau\lambda(x)\}\psi_L(x), \quad (1.51a)$$

$$l'_R(x) \rightarrow (l'_R(x))' = l'_R(x) \quad (1.51b)$$

$$v'_R(x) \rightarrow (l'_R(x))' = l'_R(x) \quad (1.51c)$$

$$A_\alpha(x) \rightarrow A'_\alpha(x) = A_\alpha(x) + \frac{1}{g}\partial_\alpha\lambda(x) - \lambda(x) \times A_\alpha(x) \quad (1.51d)$$

(where the $\lambda(x)$ are arbitrary real functions of x ($i = 1, 2, 3$), and where A_α^i is a triplet of vector fields). We assume the interaction of vector bosons to be minimal. Such an interaction is introduced via the substitution (see sec.(1.5))

$$\partial_\alpha\psi_{lL} \rightarrow \left(\partial_\alpha - ig\frac{1}{2}\tau A_\alpha\right)\psi_{lL} \quad (1.52)$$

(g is the dimensionless constant). from equations (1.50) and (1.52) we get the interaction Lagrangian of leptons and vector bosons as

$$\mathfrak{L}_i = ig\mathbf{j}_\alpha A_\alpha \quad (1.53)$$

where

$$\mathbf{j}_\alpha = \sum_l \bar{\psi}_{lL}\gamma_\alpha\frac{1}{2}\tau\psi_{lL} \quad (1.54)$$

From (1.53) we can single out the interaction of leptons with charged vector bosons:

$$\mathfrak{L}_i = \left(\frac{ig}{2\sqrt{2}}j_\alpha^{(+)}W_\alpha + h.c.\right) + igj_\alpha^3A_\alpha^3 \quad (1.55)$$

where

$$W_\alpha = \frac{1}{\sqrt{2}} (A_\alpha^1 - iA_\alpha^2) = \frac{1}{\sqrt{2}} A_\alpha^{1-i2} \quad (1.56)$$

is the field of charged vector bosons and

$$j_\alpha^{(+)} = 2j_\alpha^{1+i2} = 2 \sum_i \bar{\psi}_{iL} \gamma_\alpha \tau_+ \psi_{iL} = \sum_l \bar{v}_l' \gamma_\alpha (1 + \gamma_5) l' \quad (1.57)$$

is the charged current. therefore, the interaction Lagrangian (1.55) which follows from the local gauge invariance does contain the term (1.44) describing the interaction of leptons with a charged intermediate boson.

The second term in the Lagrangian (1.55) describes the interaction of neutrinos and charged leptons with the neutral vector boson:

$$\mathcal{L}'_i = ig \frac{1}{4} \sum_l (\bar{v}_l' \gamma_\alpha (1 + \gamma_5) v_l' - \bar{l}' \gamma_\alpha (1 + \gamma_5) l') A_\alpha^3 \quad (1.58)$$

The *GWS* theory is a unified theory of weak and electromagnetic interaction. Obviously, the interaction (1.58) is not electromagnetic interaction. For a unification of weak and electromagnetic interactions it is necessary, therefore, to require the invariance of the Lagrangian of the system with respect to a larger group than the local $SU(2)$. The simplest possibility is the group $SU(2) \times U(1)$ which makes the basis of the *GWS* theory.

To construct the locally $SU(2) \times U(1)$ invariant Lagrangian we perform in equation (1.50) the minimal substitution (section (1.5))

$$\partial_\alpha \psi_{lL} \rightarrow \left(\partial_\alpha - ig \frac{1}{2} \tau A_\alpha - ig' \frac{1}{2} y_L B_\alpha \right) \psi_{lL}, \quad (1.59a)$$

$$\partial_\alpha l'_R \rightarrow \left(\partial_\alpha - ig' \frac{1}{2} y_R^{(-1)} B_\alpha \right) l'_R, \quad (1.59b)$$

$$\partial_\alpha v'_{lR} \rightarrow \left(\partial_\alpha - ig' \frac{1}{2} y_R^{(0)} B_\alpha \right) v'_{lR}, \quad (1.59c)$$

where A_α is a triplet of gauge fields with respect to the group $SU(2)$, B_α is the gauge field associated with the symmetry group $U(1)$ and the y constants are the corresponding hypercharge. The complete gauge invariant Lagrangian of leptons and vector bosons consequently becomes

$$\begin{aligned} \mathfrak{L} = & - \sum_l \bar{\psi}_{lL} \gamma_\alpha \left(\partial_\alpha - ig \frac{1}{2} \tau A_\alpha - ig' y_L B_\alpha \right) \bar{\psi}_{lL} - \sum_l \bar{l}'_R \gamma_\alpha \left(\partial_\alpha - ig' \frac{1}{2} y_R^{(-1)} B_\alpha \right) l'_R \\ & - \sum_l \bar{v}'_{lR} \gamma_\alpha \left(\partial_\alpha - ig' \frac{1}{2} y_R^{(0)} B_\alpha \right) v'_{lR} - \frac{1}{4} \mathbf{F}_{\alpha\beta} \mathbf{F}_{\alpha\beta} - \frac{1}{4} F_{\alpha\beta} F_{\alpha\beta} \end{aligned} \quad (1.60)$$

where

$$\mathbf{F}_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha + g A_\alpha \times A_\beta \quad (1.61a)$$

$$F_{\alpha\beta} = \partial_\alpha B_\beta - \partial_\beta B_\alpha \quad (1.61b)$$

The interaction Lagrangian of leptons and vector bosons, which follows from equation (1.60), can be written as

$$\mathfrak{L}_i = ig \mathbf{j}_\alpha \mathbf{A}_\alpha + ig' \frac{1}{2} j_\alpha^y B_\alpha \quad (1.62)$$

The current j_α is given by (1.54), and

$$j_\alpha^y = \sum_l y_L \bar{\psi}_{lL} \gamma_\alpha \psi_{lL} + \sum_l y_R^{(-1)} \bar{l}'_R \gamma_\alpha l'_R + \sum_l y_R^{(0)} \bar{v}'_R \gamma_\alpha v'_R \quad (1.63)$$

The $U(1)$ invariance does not impose any constraints on the coupling constants between the leptons and the field B_α (see the discussion at the end of sec. (1.5)). This freedom in the choice of the coupling constants for the $U(1)$ gauge group can then be used to unify weak and electromagnetic interactions.

We will choose y_L , $y_R^{(-1)}$ and $y_R^{(0)}$ so as to satisfy the Gell-Mann–Nishijima relation

$$Q = I_3 + \frac{1}{2}y \quad (1.64)$$

Here Q is the electric charge in units of the proton charge, I_3 is the third component of the weak isospin. It follows that y_L equals the sum of the charges of the “upper” and “lower” components of the doublet ψ_{lL}

$$y_L = -1 \quad (1.65)$$

correspondingly, the weak hypercharges of the right-handed singlets l'_R and v'_R are equal to:

$$y_R^{(-1)} = -2, \quad y_R^{(0)} = 0 \quad (1.66)$$

respectively. With the help of equations (1.64)-(1.66) it is easy to check that

$$\frac{1}{2}j_\alpha^{(y)} = j_\alpha^{em} - j_\alpha^3 \quad (1.67)$$

where

$$j_\alpha^{em} = \sum_l (-1) \bar{l}' \gamma_\alpha l' \quad (1.68)$$

is the electromagnetic current of leptons and where j_α^3 is the third component of the isovector $\mathbf{j}_\alpha$.

Using equation (1.67) the interaction Lagrangian (1.62) can be written as

$$\mathfrak{L} = \left(\frac{ig}{2\sqrt{2}} j_\alpha^{(+)} W_\alpha + h.c. \right) + \mathfrak{L}_i^0 \quad (1.69)$$

where

$$\mathfrak{L}_i^0 = igj_\alpha^3 A_\alpha^3 + ig' (j_\alpha^{em} - j_\alpha^3) B_\alpha \quad (1.70)$$

is the interaction Lagrangian of the leptons and the neutral vector bosons.

To single out the electromagnetic interaction from equation (1.70), we rewrite this expression as

$$\mathfrak{L}_i^0 = i\sqrt{g^2 + g'^2} j_\alpha^3 \left(\frac{g}{\sqrt{g^2 + g'^2}} A_\alpha^3 - \frac{g'}{\sqrt{g^2 + g'^2}} B_\alpha \right) + ig' j_\alpha^{em} B_\alpha \quad (1.71)$$

Instead of the fields A_α^3 and B_α we introduce the field

$$Z_0 = \frac{g}{\sqrt{g^2 + g'^2}} A_\alpha^3 - \frac{g'}{\sqrt{g^2 + g'^2}} B_\alpha \quad (1.72)$$

and the field

$$A_\alpha = \frac{g'}{\sqrt{g^2 + g'^2}} A_\alpha^3 - \frac{g}{\sqrt{g^2 + g'^2}} B_\alpha \quad (1.73)$$

orthogonal to Z_α . Elementary algebra implies that the field A_α is coupled only to j_α^{em} , while the field Z_α is coupled both with the current j_α^3 and j_α^{em} . This means that the expression (1.71) contains the Lagrangian of the electromagnetic interactions and that A_α is the electromagnetic field.

Indeed, we have

$$\mathfrak{L}_i^0 = i \frac{1}{2} \sqrt{g^2 + g'^2} j_\alpha^0 Z_0 + i \frac{gg'}{\sqrt{g^2 + g'^2}} j_\alpha^{em} A_\alpha \quad (1.74)$$

where

$$j_\alpha^0 = 2 \left(j_\alpha^3 - \frac{g'}{g^2 + g'^2} j_\alpha^{em} \right) \quad (1.75)$$

If the coupling constants g and g' are related to the charge of the proton as

$$\frac{gg'}{\sqrt{g^2 + g'^2}} = e \quad (1.76)$$

the second term $ie j_\alpha^{em} A_\alpha$ in expression (1.74) becomes the interaction Lagrangian between leptons and photons.

Thus there are four vector bosons fields associated with the gauge $SU(2) \times U(1)$ group. Two fields correspond to charged vector bosons (W^+ and W^-) and two fields correspond to neutral ones. One neutral field is identified with the electromagnetic field, the other is the field of the neutral intermediate boson.

Consequently, the unification of weak and electromagnetic interactions based on the group $SU(2) \times U(1)$ is possible provided that not only charged vector bosons and charged currents but also neutral vector bosons and neutral currents, do exist.

Now we will continue our constructing of the unified electroweak theory of GPS . The Weinberg angle θ_W is introduced as follows:

$$\tan \theta_W = \frac{g'}{g} \quad (1.77)$$

For the neutral current j_α^0 we get

$$j_\alpha^0 = 2j_\alpha^3 - 2 \sin^2 \theta_W j_\alpha^{em} \quad (1.78)$$

and the relation (1.76) turns into

$$g = \frac{e}{\sin \theta_W} \quad (1.79)$$

the complete interaction Lagrangian of leptons and gauge bosons can be rewritten with the help of equations (1.69), (1.74) and (1.77) as

$$\mathfrak{L}_i = \left(\frac{ig}{2\sqrt{2}} j_\alpha^{(+)} W_\alpha + h.c \right) + i \frac{g}{2 \cos \theta_W} j_\alpha^0 Z_\alpha + ie j_\alpha^{em} A_\alpha \quad (1.80)$$

The structure of the neutral current in the GWS theory is determined by the unifying weak and electromagnetic interactions. the first term in equation (1.78) is the third component of the isovector, whose “plus-component” is identified with the charged weak current. the parameter $\sin^2 \theta_W$ is thus the only parameter which enters the expression for the neutral current. Its value can be determined from the data on the neutral current induced processes.

The theory we have considered so far satisfies the requirements of a local $SU(2) \times U(1)$ gauge invariance. Mass terms of the vector boson fields can not be introduced into the Lagrangian of such a theory. It is also obvious that the $SU(2) \times U(1)$ invariance with left-handed fields in doublets ψ_{lL} and right-handed fields l_R in singlets also forbids the introduction of lepton mass terms into the Lagrangian.

In the standard electroweak theory the Lagrangian mass terms of the both the vector boson and fermion fields are introduced by the Higgs mechanism of spontaneous breakdown of the gauge symmetry (see sec. (1.6)). The theory is built up so that, at the beginning, the complete Lagrangian, including the Higgs sector, is locally $SU(2) \times U(1)$ invariant. It is then necessary to assume that the Higgs fields transformed according to a definite (non-trivial) representation of the gauge group. Further, due to the spontaneous breakdown of the gauge invariance charged (W^+ and W^-) as well as neutral Z^0 intermediate bosons have to acquire masses. That is, three Goldstone degrees of freedom of Higgs field can transform at the spontaneous breakdown of the gauge invariance into the additional degrees of freedom of vector fields (three masses). Thus, we are forced

to assume that the Higgs fields form at least doublet. It is this “minimal” assumption which is at the bases of the GBS theory.

hence we assume that the Higgs fields forma doublet of the $SU(2)$ group

$$\phi = \begin{pmatrix} \phi_+ \\ \phi_0 \end{pmatrix} \quad (1.81)$$

where the complex function ϕ_+ , ϕ_0 are the fields of the charged and neutral bosons, respectively. Weak hypercharge of the doublet (1.81) is defined so as to fulfill the Gell-Mann–Nishijima relation (1.64). We have

$$y_\phi = 1 \quad (1.82)$$

The Lagrangian of the Higgs field $\phi(x)$ is given as (see sec.(1.6))

$$\mathcal{L}^0 = -\partial_\alpha \phi^+ \partial_\alpha \phi - V(\phi^+ \phi). \quad (1.83)$$

Here

$$V(\phi^+ \phi) = -\mu^2 \phi^+ \phi + \lambda (\phi^+ \phi)^2 = \lambda \left(\phi^+ \phi - \frac{\mu^2}{2\lambda} \right)^2 - \frac{\mu^4}{4\lambda} \quad (1.84)$$

where μ^2 and λ are positive constants.

Taking into account (1.82), we get from (1.83) by the standard substitution

$$\partial_\alpha \phi \rightarrow \left(\partial_\alpha - ig \frac{1}{2} \tau A_\alpha - ig' \frac{1}{2} B_\alpha \right) \phi$$

the Lagrangian

$$\mathcal{L} = - \left[\partial_\alpha \phi^+ + \phi^+ \left(ig \frac{1}{2} \tau A_\alpha + ig' \frac{1}{2} B_\alpha \right) \right] \left[\partial_\alpha \phi + \left(ig \frac{1}{2} \tau A_\alpha + ig' \frac{1}{2} B_\alpha \right) \phi \right] - V(\phi^+ \phi) \quad (1.85)$$

is invariant with respect to the gauge group $SU(2) \times U(1)$. It is obvious from (1.84) that the potential $V(\phi^+ \phi)$ is minimal for

$$(\phi^+ \phi)_0 = \frac{\mu^2}{2\lambda} = \frac{v^2}{2} \quad (1.86)$$

For the minimal (vacuum) value of ϕ we choose

$$\phi_{vac.} = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix} \quad (1.87)$$

Further, the doublet ϕ can always be written in the form

$$\phi(x) = \exp \left\{ i \frac{1}{2} \tau \frac{\theta(x)}{v} \right\} \begin{pmatrix} 0 \\ \frac{\{v + \chi(x)\}}{\sqrt{2}} \end{pmatrix} \quad (1.88)$$

where $\theta(x)$ and $\chi(x)$ are real functions. Finally, the functions $\theta(x)$, which, correspond to the “would be” Goldstone bosons can always be eliminated owing to the gauge invariance of the Lagrangian (1.85) by appropriately fixing the gauge (the so called unitary gauge). Thus we have

$$\phi(x) = \begin{pmatrix} 0 \\ \frac{\{v + \chi(x)\}}{\sqrt{2}} \end{pmatrix} \quad (1.89)$$

Let us substitute (1.89) into (1.85). Taking into account that

$$(\boldsymbol{\tau} \mathbf{A}_\alpha) (\boldsymbol{\tau} \mathbf{A}_\alpha) = 2W_\alpha \overline{W}_\alpha + A_\alpha^3 A_\alpha^3,$$

$$\phi^+ (\boldsymbol{\tau} \mathbf{A}_\alpha) B_\alpha \phi = -A_\alpha^3 B_\alpha \frac{1}{2} (v + \chi)^2$$

we get

$$\mathfrak{L} = \frac{1}{2} \partial_\alpha \chi \partial_\alpha \chi - \frac{1}{2} (v + \chi)^2 \left[\frac{1}{4} g^2 2W_\alpha \overline{W}_\alpha + \frac{1}{4} (g^2 + g'^2) Z_\alpha Z_\alpha \right] - \frac{1}{4} \lambda \chi^2 (\chi + 2v)^2 \quad (1.90)$$

Here

$$W_\alpha = \frac{A_\alpha^{1-i2}}{\sqrt{2}} \quad \text{and} \quad \overline{W}_\alpha = \frac{A_\alpha^{1+i2}}{\sqrt{2}}$$

are the fields of the charged vector bosons and Z_α is the field of the neutral vector bosons.

As a results of the spontaneous breakdown of the symmetry, mass term for the intermediate bosons have emerged from in the Lagrangian

$$\mathfrak{L}_m = -m_W^2 W_\alpha \overline{W}_\alpha - \frac{1}{2} m_Z^2 Z_\alpha Z_\alpha, \quad (1.91)$$

where

$$m_W^2 = \frac{1}{4} g^2 v^2, \quad m_Z^2 = \frac{1}{4} (g^2 + g'^2) v^2. \quad (1.92)$$

Symmetry was broken in such a way that the photon remained a massless particle.

The function $\chi(x)$ is a field of neutral scalar particles (the so called Higgs particles).

It follows from (1.90) that their mass is equal to

$$m_\chi = \sqrt{2\lambda}v = \sqrt{2}\mu \quad (1.93)$$

note that the Lagrangian (1.90) contains also a term describing the interaction of the Higgs particles with the intermediate bosons.

We find from (1.77) and (1.92) that the mass squared of the Z boson is related to that of the W boson and the parameter $\cos^2 \theta_W$ by

$$m_Z^2 = \frac{m_W^2}{\cos^2 \theta_W}. \quad (1.94)$$

It should be stressed that this relation is satisfied only if the Higgs fields form doublets. In the case of higher Higgs multiplets no relation between masses of neutral and charged intermediate bosons does exist.

It follows from (1.45) and (1.92) that

$$v = \frac{1}{(\sqrt{2}G_F)^{\frac{1}{2}}} \quad (1.95)$$

Therefore, the theory enables us to calculate the parameter v . Substituting the numerical value of G_F :

$$G_F = 1.1664 \times 10^{-5} GeV,$$

we find

$$v = 246.2 GeV. \quad (1.96)$$

Also, it follows from (1.45) and (1.79) that

$$m_W = \left(\frac{\pi\alpha}{\sqrt{2}G_F} \right)^{\frac{1}{2}} \frac{1}{\sin \theta_W}. \quad (1.97)$$

the value of the parameter $\sin \theta_W$ is determined from experimental data on neutral currents induced processes. therefore, the theory enables us to predict the value of the W boson mass.

From the analysis of the world data on deep inelastic processes, one could deduce the value

$$\sin^2 \theta_W = 0.2315. \quad (1.98)$$

With the above value of $\sin^2 \theta_W$, the masses of charged and neutral intermediate bosons using (1.97) and (1.94) turn out to be

$$m_W = 80.330 GeV \quad (1.99)$$

$$m_Z = 91.187 GeV \quad (1.100)$$

CHAPTER 2

SUPERSYMMETRY

2.1 Motivation *for* Supersymmetry

Ever since its discovery in the early seventies, supersymmetry has been the focus of considerable attention. Although no compelling supersymmetric model has yet emerged, and in spite of the fact that there is no experimental evidence for supersymmetry, its remarkable theoretical properties have provided sufficient motivation for its study.

Supersymmetry, is a novel symmetry that interrelates bosons and fermions, thereby providing a new level of synthesis. It is the most general (known) symmetry of the $S - matrix$ consistent with *Poincare'* invariance. Supersymmetry leads to an improvement (and sometimes even to elimination of divergencies that occur in Quantum Field Theory (QFT), in particular, quadratic divergencies are absent); these feature will play an important role in our subsequent discussions. Since two successive supersymmetry transformation involve a space-time translation, local supersymmetry theories (and past experience shows that nature prefers local supersymmetry!) necessarily include gravitation with the gauge fermion (the *gravitino*) being the supersymmetric partner of the *graviton*. Further, because supersymmetric QFT's exhibit a better ultraviolet behavior, they may provide a hope of eventually obtaining a consistent quantum theory that in-

clude gravitation. Finally, supersymmetry is an essential ingredient in the construction of the most recent candidate for a *theory of everything*, *TOE*, the *SUPERSTRING*.

At this point, one may ask why any extension of the Standard Model (SM) needs to be considered [7]. After all, the *GWS* theory seems for all known electromagnetic and weak phenomena, and quantum chromodynamics (*QCD*) is generally accepted to be the theory of strong interactions. Indeed, it appears that all experiments are consistent with a gauge theory based on $SU(3)_c \times SU(2)_L \times U(1)_Y$, with the $SU(2)_L \times U(1)_Y$ being spontaneously broken to $U(1)_{em}$.

In the *GWS* model, the spontaneous breakdown is brought about the introduction of an elementary scalar field. This leads to the prediction of (at least) an additional spin-zero particle, the Higgs boson. With the discovery of the $W^\pm$ and Z^0 bosons at the *CERN* $p\bar{p}$ collider, and the top quark at the *FERMILAB*, only the Higgs boson remains to be discovered to complete the particle content of the *GWS* electroweak theory. It should be pointed out that no elementary spin-zero particles have ever been found. In fact, the *ad hoc* introduction of these considered by many theorists to be an unpleasant feature of the standard model.

The problem is the instability of the scalar particles masses under radiative corrections. For example, one-loop radiative corrections to these diverge quadratically, leading to corrections of the form:

$$\delta m^2 = O(\alpha/\pi)\Lambda^2 \tag{2.1}$$

where Λ is a cut-off parameter representing the scale of the theory and $\alpha = e^2/hc \approx \frac{1}{137}$ is the fine structure constant. The parameter Λ may be the *Grand Unified Theory* (*GUT*) scale $O(10^{15}GeV/c^2)$, or the *Plank scale* $O(10^{19}GeV/c^2)$ if we believe there is no new physics all the way up to the scale associated with quantum gravity. On the other hand, we know that for the scalar self-couplings to be sensibly treated within perturbation theory, the scalar mass

$$m^2 \leq O(m_W^2/\alpha) \sim 1TeV/c^2 \quad (2.2)$$

In other words, either the Higgs sector is strongly interacting, or $\delta m^2 \gg m^2$. Such theories (where $\delta m^2 \gg m^2$) have been technically referred to as “unnatural”, because the parameters have to be tuned with unusual precision in order to preserve the lightness of the Higgs mass compared to the GUT scale $\Lambda \sim O(1TeV/c^2)$

One possible solution to the problem of naturalness is to imagine that the quarks, leptons and gauge bosons are all composites with associated scale $\Lambda \sim O(1TeV/c^2)$. While these solve the problem at present energies, it does not really represent a solution as we could ask the same questions of any underlying theory. Moreover, it would be difficult to understand why the gauge principle seems to work so well at least up to this point.

A different approach would be to eliminate the fundamental scalars and imagine that the Higgs is a composite of new fermions bound by a new force -*the technicolor force*- that becomes strong at a scale of $O(1TeV/c^2)$. While this is an appealing idea, and while the composite Higgs boson can indeed lead to masses for the $W^\pm$ and Z^0 , it

does not account for quark and lepton masses. This led to the introduction of the yet another interaction, the extended technicolor. This appears to have phenomenological problems, particularly with flavor-changing neutral currents (FCNC). Although that it appears there is no reason for the technicolor idea not to work, it seems fair to say that no viable model has yet emerged.

Supersymmetry provides yet another to the naturalness question. this may be simply understood by recalling that fermion and masses are protected from large radiative corrections by chiral symmetry. The analogue for equation (2.1) is

$$\delta m_f = O(\alpha/\pi) m_t \ln(\Lambda/m_t) \quad (2.3)$$

Thus, massless fermions do not acquire masses via radiative corrections. This is a manifestation of the chiral symmetry. The naturalness problem arises because, unlike the case of fermions, there is no symmetry to keep massless scalars from acquiring large masses via radiative corrections.

In practice, at the one-loop level, this works because boson and fermion loops both enter the scalar correction, but with a relative minus sign. For supersymmetric theories, equation (2.1) takes the form

$$\delta m^2 \approx O(\alpha/\pi)\Lambda^2 - O(\alpha/\pi)\Lambda^2 = 0 \quad (2.4)$$

Exact cancellation requires that the bosons and fermions enter with the same quantum numbers (this is ensured by supersymmetry) so that their couplings are the same

except for supersymmetry Clebsch-Gordon factors, and also that they have the same masses. Spontaneous breaking of supersymmetry as for any other symmetry, maintains a relation between the couplings but breaks the mass relations, and equation (2.4) takes the form

$$\delta m^2 \approx O(\alpha/\pi) |m_B^2 - m_f^2| \quad (2.5)$$

We see from expression (1.2) and (1.5) that for supersymmetry to solve the naturalness problem,

$$|m_B^2 - m_f^2| \leq O(1TeV/c^2)^2 \quad (2.6)$$

where m_B^2 (m_f^2) is the boson (fermion) to have masses $\leq O(1TeV/c^2)$ and hope that these may show up at future LEP energies.

We emphasize that not one of all these reasons for considering supersymmetry, no matter how compelling it may appear, requires any particular mass scale for supersymmetric particles (sparticles). It is only if we require supersymmetry to address the naturalness question that the mass scale is fixed. We note that the mass scale does address the question " why is the scalar mass 12 (16) order of magnitude smaller than the *GUT* (*Planck*) scale in the first place?". But in supersymmetric theories, once this value has been set, either "by hand" or by any other mechanism, radiative corrections preserve this hierarchy of scales.

2.2 Rules *of* Supersymmetry

We are here to establish some of the rules of supersymmetry treatment [8].

First and foremost we postulate the *existence* of supersymmetry between fermions and bosons which should underlay the laws of physics. No experimental observation has yet revealed particles or forces which manifestly show such a symmetry except for some rare events [9]. therefore the development of a theory based on supersymmetry requires an understanding not only of how the various symmetry transformations affect each other (the algebra) but also of all possible systems (*multiplets* of particles or quantum fields) on which the supersymmetry transformations can act. The symmetry operations will transform different members of a multiplet into each other. More precisely, the transformations are to be represented by linear operators acting on the vector space. (the “*representation space*”) spanned by the multiplet. Finally the theory must predict the time development of interacting physical systems. This is usually achieved by finding appropriate *Hamiltonians* or *Lagrangians*. The supersymmetry present in the physical system will manifest itself in the invariance of this *Lagrangian* - or rather its integral over all time, **the action**- if all the fields undergo their respective supersymmetry transformation. Because of the lack of experimental input, a large fraction of the research effort of supersymmetry theorists has, in fact, been devoted to the finding of, and exploration of, possible supersymmetry respecting interactions.

The theoretical framework in which to construct supersymmetric models in flat space-time is quantum field theory, and it must be pointed out that *the standard con-*

cept of quantum field theory allow for supersymmetry without any further assumptions. This introduction of supersymmetry is not a revolution in the way one views physics. It is an additional symmetry that in otherwise “normal” field theoretical model can have. As we shall see, all that is required for a field theory to be supersymmetric is that it contains specified types and numbers of fields in interactions with each other and that the various interaction strengths and particle masses have properly related events. As an example, consider the $SU(3)$ gauge theory of gluons, which can be made supersymmetric by including a massless neutral color octet of spin $\frac{1}{2}$ particles which are their own anti-particles. Such spin $\frac{1}{2}$ partners of the gluons are called “*gluinos*”. If our model contains not only gluons but also quarks, we must also add corresponding partners for them. These have spin 0 and are commonly called “*squarks*”. (Procedures like these are employed particularly in the construction of supersymmetric Grand Unified Theories *susyGUTs* or *superGUTs*.)

Before we proceed to discuss the ingredients of supersymmetric models, we must address the question of the *Fermi – Bose* matter-force duality. After all, the wave particle duality of quantum mechanics and the subsequent question of the “exchange particle” in perturbative quantum field theory seemed to have abolished that distinction for good. The recent triumph progress of gauge theories has, however, reintroduced it. forces are mediated by gauge potentials, i.e., from spin $\frac{1}{2}$ *fermions*. The Higgs particles, necessary to mediate the needed spontaneous breakdown of the gauge invariances (more about this later), play a somewhat intermediate role. They must have zero spin and

are thus bosons, but they are not directly related to any of the forces. Purists hope to see them arise as bound states of the fermions (condensates). Supersymmetric theories, and particularly supergravity theories, “unite” fermions and bosons into multiplets and left the basic distinction between matter and interaction. The gluinos, for example, are thought of as carriers of the strong force as much as the gluons, except that as fermions they must obey an exclusion principle and thus will never conspire to form a coherent, measurable potential. The distinction between forces and matter becomes phenomenological: bosons - and particularly massless ones - manifest themselves as forces because they can build up coherent classical fields; fermions are seen as matter because no two identical ones can occupy the same point in space.

For some time it was thought that supersymmetry which would naturally relate forces and fermionic matter would be in conflict with field theory. The progress in understanding elementary particles through the $SU(3)$ classification of the “eight-fold way” (a global symmetry) had led to attempts to find a unifying symmetry which would directly relate to each other several of the $SU(3)$ multiplets (baryon octet, decuplet, etc.) even if these had different spins. The failure of attempts to make those “spin symmetry” relativistically covariant led to the formulation of a series of no-go theorems, culminating in 1967 in a paper by *Coleman* and *Mandula* which was widely understood to show that it is impossible, within the theoretical framework of relativistic field theory, to unify space-time symmetry with internal symmetries. More precisely, say that the charge operators whose eigenvalues represent the “internal” quantum numbers such as electric

charge, isospin, hypercharge, etc., must be transitionally invariant. This means that these operators commute with the energy, the momentum operators. indeed the only symmetry generators which transfer at all under both translations and rotations are those of the *Lorentz* transformations themselves (rotations and transformations to coordinate systems which move with constant velocity). The generators of internal symmetries can not relate eigenstates with different eigenvalues m^2 and $l(l+1)\hbar^2$ of the mass and spin operators. This means that irreducible multiplets of symmetry groups can not contain particles of different mass or of different spin. This no-go theorem, seemed to rule out exactly the sort of unity which was sought. One of the assumptions made in Coleman and Mandula's proof did, however, turn out to be unnecessary: they had admitted only those symmetry transformations which form *Lie* group with real parameters. Examples of such symmetries are space rotations with the *Euler* angles as parameters and the phase transformations of electrodynamics with a real phase angel θ about which we talked earlier. The charge operators associated with such Lie groups of symmetry transformations (their generators) obey well-defined *commutation relations* with each other. Perhaps the best-known example is the set of commutators $L_x L_y - L_y L_x \equiv [L_x, L_y] = i\hbar L_z$, for the angular momentum operators which generate spatial rotations.

Different spins in the same multiplet are allowed if one includes symmetry operations whose generators obey *anticommutation relations* of the form $AB + BA \equiv \{A, B\} = C$. This was first proposed in 1971 by Gol'fand and Likhtman, and followed up by Volkov

and Akulov who arrived at what we now call a non-linear realization of supersymmetry. Their model was not renormalizable. In 1973, Wess and Zumino presented a renormalizable theoretical model of a spin $\frac{1}{2}$ particle in interaction with two spin 0 particles where the particles are related by symmetry transformations, and therefore “sit” in the same multiplet. The limitation of the Coleman-Mandula no-go theorem had been avoided by the introduction of a fermionic symmetry operator which carried a spin $\frac{1}{2}$, and thus when acting on a state of spin j resulted in a linear combination of states with spin $j + \frac{1}{2}$ and $j - \frac{1}{2}$. Such operators must and do observe anticommutator relations with each other. They do not generate Lie groups and are therefore not ruled out by the Coleman-Mandula no-go theorem. In the light of this discovery, Haag, Lopuszanski, and Sohnius extended the results of Coleman-Mandula to include symmetry operations which obey Fermi-statistics. They proved that in the context of relativistic field theory the only model which can lead to a solution of the unification problems are supersymmetric theories, and space-time and internal symmetries can only be related to each other by fermionic symmetry operators Q of spin $\frac{1}{2}$ (*not $\frac{3}{2}$ or higher*) whose properties are either exactly those of the Wess-Zumino model or are at least closely related to them. Only in the presence of supersymmetry can multiplets contain particles of different spin, such as the graviton and the photon.

2.3 Essentials of Supersymmetry Algebra

Supersymmetry transformations are generated by quantum operators Q which change fermionic states to bosonic ones and vice versa,

$$Q|fermion\rangle = |boson\rangle; \quad Q|boson\rangle = |fermion\rangle. \quad (2.7)$$

Which particular bosons and fermions are related to each other by the operation of some such Q ?, how many Q 's there are? and which properties other than statistics of the states are changed by that operation depends on the supersymmetric model under study. There are, however, a number of properties which are common to the Q 's in all supersymmetric models.

By definition, the Q 's change the statistics and hence the spin of the state. Spin is related to behavior under spatial rotations, and thus supersymmetry is- in some sense- a space-time symmetry. Normally, and particularly so in models of “extended supersymmetry” (supergravity is being one example), the Q 's also affect the internal quantum numbers of the states. It is this property of combining internal with space-time behavior that makes supersymmetric field theories interesting in the attempts to unify all fundamental interactions.

As a simple illustration of the non-trivial space-time properties of the Q 's, consider the following. Because fermions and bosons behave differently under rotations, the Q can not be invariant under such rotations. We can, for example, apply the unitary operator $U(U^{-1}U = 1)$ which, in Hilbert space, represents a rotation in configuration

space by 360^0 around some axis. Since fermionic states pick up a minus sign when rotated through 360^0 and bosonic states do not, we have

$$U |fermion\rangle = -|fermion\rangle, \quad U |boson\rangle = |boson\rangle, \quad (2.8)$$

then from equation (2.1) we get

$$\begin{aligned} Q |boson\rangle &= |fermion\rangle = -U |fermion\rangle = -UQ |boson\rangle = -UQU^{-1}U |boson\rangle \\ &= -UQU^{-1} |boson\rangle, \end{aligned} \quad (2.9)$$

$$\begin{aligned} Q |fermion\rangle &= |boson\rangle = U |boson\rangle = UQ |fermion\rangle = UQU^{-1}U |fermion\rangle \\ &= -UQU^{-1} |fermion\rangle, \end{aligned} \quad (2.10)$$

and since all fermionic and bosonic states, taken together, form a basis in the Hilbert space, we easily see that we *must* have

$$UQU^{-1} = -Q \quad (2.11)$$

The rotated supersymmetry generator picks up a minus sign, just as a fermionic state does. One can extend this analysis and show that the behavior of the Q 's under any Lorentz transformation -not only under rotations by 360^0 - is precisely that of a *spinor operator*. More technically speaking, the Q 's transform like tensor operators of spin $\frac{1}{2}$

and, in particular, they do not commute with Lorentz transformations followed by a supersymmetry transformation is different from that when the order of the transformation is reserved.

It is not easy to illustrate, but it is nevertheless true that, on the other hand, the Q 's are *invariant under transformation*. It does not matter whether we translate the coordinate system before or after we perform a supersymmetry transformation. In technical terms, this means that we have a vanishing commutator of Q with the energy of momentum operator E and $\mathbf{P}$, which generates space-time translations,

$$[Q, E] = [Q, \mathbf{P}] = 0 \tag{2.12}$$

The structure of a set of symmetry operations is determined by the result of two subsequent operations. For continuous symmetries like space rotations or supersymmetry, this structure is best described by the commutators of the generators, such as the ones given above for the angular momentum operators. The commutator structure of the Q 's with themselves can best be examined by if they are viewed as products of operators which annihilate fermions and create bosons instead, or vice versa. It can be shown that the canonical quantization rules for creation and annihilation operators of particles (and in particular the *anticommutator* rules for fermions, which reflect Pauli's exclusion principle) lead to the results that it is the anti commutator of two Q 's, not the commutator, which is again a supersymmetry generator, albeit one of bosonic nature.

Let us consider the anticommutator of some Q with its Hermitian adjoint Q^+ . As

spinor components, the Q 's are in general not Hermitian, but $\{Q, Q^+\} \equiv QQ^+ + Q^+Q$ is a Hermitian operator with positive definite eigenvalues;

$$\langle \dots | QQ^+ | \dots \rangle + \langle \dots | Q^+Q | \dots \rangle = |Q | \dots \rangle|^2 + |Q^2 | \dots \rangle|^2 \geq 0. \quad (2.13)$$

This can only be zero for all states $| \dots \rangle$ if $Q = 0$. A more detailed investigation will show that $\{Q, Q^+\}$ must be a linear combination of the energy and momentum operators;

$$\{Q, Q^+\} = \alpha E + \beta \mathbf{P} \quad (2.14)$$

This relation between the anticommutator of two generators of supersymmetry transformations on the one hand and the generators of *space – time translations* (namely energy and momentum) on the other, is central to the entire field of supersymmetry and supergravity. It means that the subsequent operations of two finite supersymmetry transformations will include translations in space and time of the states on which they operate.

There is a further important consequence of the form of equation (2.14). When summing this equation over all supersymmetry generators, we find that $\beta \mathbf{P}$ terms cancel while the αE terms add up, so that

$$\sum_{all Q} \{Q, Q^+\} \propto E. \quad (2.15)$$

Depending on the sign of the proportionality factor, the spectrum for the energy

would have to be either ≥ 0 or ≤ 0 because of the inequality (2.13). For a physical sensible theory with energies bounded from below but not from above, the proportionality factor will therefore be positive.

The equation (2.12) to (2.15) are crucial properties of the supersymmetry generators, and many of the most important features of supersymmetric theories, whether in flat or curved space-time, can be derived from them.

One such feature is the *positivity of energy*, as can be seen from equation (2.15) in conjunction with (2.13), *the spectrum of the energy operator E (the Hamiltonian) in a theory with supersymmetry contains no negative eigenvalues*. We denote the state (or the family of states) with the lowest energy by $|0\rangle$ and call it the *vacuum*. The vacuum will have zero energy

$$E |0\rangle = 0 \tag{2.16a}$$

if and only if

$$Q |0\rangle = 0 \quad \text{and} \quad Q^+ |0\rangle = 0 \quad \text{for all } Q \tag{2.16b}$$

Any state whose energy is not zero, e.g. any one-particle state, can not be invariant under supersymmetry transformations. This means that there must be one (or more) superpartner state $Q |1\rangle$ or $Q^+ |1\rangle$ for every one-particle state $|1\rangle$. thus: *each supermultiplet must contain at least one boson and one fermion whose spins differ by $\frac{1}{2}$* . A supermultiplet is a set of quantum states (or, in different context, of quantum fields)

which can be transformed into one another by one or more supersymmetry transformations. This is exactly analogous to the concept of “multiplet” known from atomic, nuclear and elementary particle physics where e.g., the proton and the neutron form an isospin doublet and can be transformed into each other by an isospin rotation.

The translational invariance of Q , expressed by equation (2.12) implies that Q does not change energy and momentum and that therefore: *all states in a multiplet of unbroken symmetry have the same mass.* Experiments do not show elementary particles to be accompanied by superpartners. with different spins but identical mass. Thus, if supersymmetry is fundamental to nature, it can only be realized as a *spontaneously broken symmetry.*

The term “spontaneously broken symmetry” is used when the interaction potentials in a theory, and therefore the basic dynamics, are symmetric but the states with lowest energy, the ground state or vacuum, is not. If a generator of such symmetry acts on the vacuum the result will not be zero. Perhaps the most familiar example of a spontaneously broken symmetry is the occurrence of ferromagnetism in spite of the spherical symmetry of the laws of electrodynamics. because the dynamics retain essential symmetry of the theory, states with very high energy tend to lose the memory of the asymmetry of the ground state and the “spontaneously broken symmetry” gets re-established. high energy may mean high temperatures, essentially because the statistics of the occupation of states is different from fermions and bosons.

If supersymmetry is spontaneously broken, the ground state will not be invariant

under all supersymmetric operations: $Q|0\rangle \neq 0$ or $Q^+|0\rangle \neq 0$ for some Q . From what we said above in equation (2.16), we conclude that: *supersymmetry is spontaneously broken if and only if the energy of the lowest lying state (the vacuum) is not exactly zero*. Whereas spontaneous supersymmetry breaking may lift the mass degeneracy of the supermultiplets by giving different masses to different members of the multiplet spectrum itself will remain intact. In particular, we still need “superpartners” for all known elementary particles, although these may not be superheavy or otherwise experimentally unobtainable. The superpartners carry a new quantum number (called R-charge). It has been shown that the highly desirable property of super GUTs model, mentioned in the introduction, namely that they stabilize the GUT hierarchy, is closely associated with a strict conservation law for this quantum number. If nature works that way the lightest particle with a non-zero R-charge must be stable. Whereas this particle may be so weakly interacting that it has not yet been observed, its presence in the universe could crucially and measurably influence cosmology.

As a matter of convention, fermionic superpartners of known bosons are denoted by the suffix -ino (hence “ gravitino, photino, gluino”); the bosonic superpartners of fermions are denoted by the prefixes s- (“squark, slepton”). The discovery of any such bosinos or sfermions would confirm the important prediction of superpartners which is common to all supersymmetric models. it would be a major breakthrough and would establish supersymmetry as an important property of the physics of nature rather than just an attractive hypothesis.

We have not yet specified “how much” supersymmetry there should be. Do we propose one spin $\frac{1}{2}$ photino as a partner of a physical photon, or two, or how many? Different supersymmetric models give different answers, depending on how many supersymmetric generators Q are present, as conserved charges, in the model. As already said, the Q s are spinor operators, and a spinor in four space-time dimensions must have at least four real components. The total number of Q 's must therefore be a multiple of four. A theory with minimal supersymmetry would be invariant under the transformations generated by just the four independent components of a single spinor operator Q_α with $\alpha = 1, \dots, 4$. We call this *a theory with N=1 supersymmetry*, and it would give rise to, e.g., a single uncharged massless spin $\frac{1}{2}$ photino which is its own antiparticle (a “Majorana neutrino”). If there is more supersymmetry, then there will be several spinor generators with four components each, $Q_{\alpha i}$ with $i = 1, \dots, N$, and we speak of a *theory with N-extended supersymmetry*, which will give rise to N-photinos. The fundamental relationship (2.14) between the generators of supersymmetry is now replaced by

$$\{Q_i, Q_j^+\} = \delta_{ij} (\alpha E + \beta \mathbf{P}) \quad (2.17)$$

2.4 The Algebra of N=1 supersymmetry

Coleman and Mandula (1967) showed that under very general assumptions, a Lie group that contains both the Poincare' group P and an internal symmetry group G must be just a direct product of P and G [10]. the generators of the Poincare' group

are the four-momentum $P_\mu (P_0 = E = H, P_1 = P_x, P_2 = P_y, P_3 = P_z)$, which produces space-time translations, and the antisymmetric tensors $M_{\mu\nu}$, which generates space-time rotations, that is

$$\mathbf{J} \equiv (M_{23}, M_{31}, M_{12}); \quad \mathbf{K} \equiv (M_{01}, M_{02}, M_{03}), \quad (2.18)$$

Where the regular momentum operator J_i generates space rotations about the i – axis and K_i generates Lorentz boosts along the i – axis. So, if the generators of the internal supersymmetry group G are denoted by T_a , the Coleman-Mandula theorem requires that

$$[P_\mu, T_a] = [M_{\mu\nu}, T_a] = 0 \quad (2.19)$$

This no-go theorem shows that it is impossible to mix internal and Lorentz space-time symmetries in a non-trivial way. Supersymmetry escapes this “no-go” theorem because, in addition to the generators $P_\mu, M_{\mu\nu}, T_a$ which satisfy commutation relations, it involves fermionic generators Q that satisfy anti-commutation relations. If we call the generators $P_\mu, M_{\mu\nu}, T_a$ “even”, and Q “odd”, then the supersymmetry algebra has the general structure

$$\begin{aligned} [even, even] &= even, \\ [odd, odd] &= even, \\ [even, odd] &= odd, \end{aligned} \quad (2.20)$$

The above is called *graded Lie Algebra*.

We now present the simplest form of supersymmetry algebra (N=1 supersymmetry). We introduce four generators Q_α ($\alpha = 1, \dots, 4$), which form a four-component Majorana spinor. Majorana spinors are the simplest possible type of spinor. They are self-conjugate, i.e.

$$Q = Q^c = C\overline{Q}^\dagger, \quad (2.21)$$

and hence have only half as many degrees of freedom as a Dirac spinor. Indeed, any Dirac spinor ψ may be written

$$\psi = \frac{1}{\sqrt{2}}(\psi_1 + i\psi_2), \quad (2.22)$$

where

$$\psi_1 = \frac{1}{\sqrt{2}}(\psi + \psi^c), \quad \text{and} \quad \psi_2 = -\frac{i}{\sqrt{2}}(\psi - \psi^c) \quad (2.23)$$

are two independent Majorana spinors that satisfy $\psi_i = \psi_i^c$.

Since Q_α is a spinor, it must satisfy

$$[Q_\alpha, M_{\mu\nu}] = \frac{1}{2}(\sigma_{\mu\nu})_{\alpha\beta} Q_\beta \quad (2.24)$$

This relation expresses the fact that the Q_α transform as a spinor under the rotation generated by $M_{\mu\nu}$ (recall that $\sigma_{\mu\nu}$ when sandwiched between spinors, transform as an antisymmetric tensor). The Jacobi identity of commutators

$$[[Q_\alpha, P_\mu], P_\nu] + [[P_\nu, Q_\alpha], P_\mu] + [[P_\mu, P_\nu], Q_\alpha] = 0 \quad (2.25)$$

requires that Q_α must be transnational invariant, that is

$$[Q_\alpha, P_\mu] = 0. \quad (2.26)$$

It is the remaining anticommutation relation

$$\{Q_\alpha, \overline{Q_\beta}\} = 2(\gamma^\mu)_{\alpha\beta} P_\mu \quad (2.27)$$

which we shall derive later, and which closes the algebra, that has the most interesting consequences. Clearly the anticommutator has to yield an even generator, which might be either P_μ or $M_{\mu\nu}$. A term of the form $\sigma^{\mu\nu} M_{\mu\nu}$ on the right-hand side would violate a generalized Jacobi identity involving Q_α , Q_β and P_μ and the algebra would not close. Indeed, if go back to the “no-go” theorem and allow for anticommutators as well as commutators, we find that the only allowed supersymmetries (apart from trivial extension) are those based on the graded Lie algebra defined by eqs. (2.24)-(2.27).

we choose Q_α to be Majorana spinor with four independent (real) parameters, but we could have used a Weyl spinor with two complex components equally well. In fact, we shall find it more convenient to work with a left-handed Weyl spinor ψ_α with $\alpha = 1, 2$, and the chiral representation of the Dirac matrices in which

$$\gamma = \begin{pmatrix} 0 & \sigma \\ -\sigma & 0 \end{pmatrix}, \gamma_0 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \gamma_5 = \begin{pmatrix} -I & 0 \\ 0 & I \end{pmatrix}, C\gamma_0^T = \begin{pmatrix} 0 & i\sigma_2 \\ -i\sigma_2 & 0 \end{pmatrix}. \quad (2.28)$$

Using the two component Weyl spinor ψ_α , we can construct a Majorana spinor in this chiral representation we find

$$Q = Q^c = \begin{Bmatrix} \psi \\ 0 \end{Bmatrix} + C\gamma_0^T \begin{Bmatrix} \psi^* \\ 0 \end{Bmatrix} - \begin{Bmatrix} \psi \\ -i\sigma\psi^* \end{Bmatrix}. \quad (2.29)$$

We then look for possible supersymmetric representations that contain massless particles. these should be the relevant multiplets, the particles we observe are thought to acquire their masses only as a result of spontaneous symmetry breaking. The procedure we employ is to evaluate the anticommutator (2.27) for a massless particle moving along the z-axis with $P_\mu = (E, 0, 0, E)$. On substituting (2.29) into equation (2.27), we find

$$\{\psi_a, \psi_b^+\} = 2E(1 - \sigma_3)_{ab} \quad (2.30)$$

with $a, b = 1, 2$, giving

$$\{\psi_1, \psi_2^+\} = 0, \{\psi_1, \psi_1^+\} = 0, \{\psi_2, \psi_2^+\} = 4E. \quad (2.31)$$

We see that ψ_2^+ and ψ_2 act as creation and annihilation operators.

Now, a massless particle of spin s can only have helicities $\lambda = \pm s$, so, starting from the left-handed state $|s, \lambda = -s\rangle$, which is annihilated by ψ_2 , only one new state can be formed, i.e., $\psi_2^+ |s, -s\rangle$. This describes a particle of spin $s + \frac{1}{2}$ and helicity $-(s + \frac{1}{2})$,

and by virtue of (2.26) it is also massless. Then, acting again with ψ_1^+ or with ψ_2^+ gives states with zero norm by virtue of (2.31) and $\psi_2^+\psi_2^+ = 0$ (which follows from the fermionic nature of ψ_2). So the resulting massless irreducible representation consists of just two states. Hence, the possible supersymmetric multiplets of interest to us are

Chairal multiplet	Vector (or gauge) multiplet
fermion $ \frac{1}{2}, \frac{1}{2}\rangle$	gauge boson $ 1, 1\rangle$
sfermion $ 0, 0\rangle$	gaugino $ \frac{1}{2}, \frac{1}{2}\rangle$

To maintain *CPT* invariance we must add the antiparticle states that have opposite helicity, thus giving a total of four states, $|s \pm \frac{1}{2}, \pm(s + \frac{1}{2})\rangle, |s, \pm s\rangle$, in each multiplet.

All the particles in such multiplets carry the same gauge quantum numbers. For this reason, the know fermions (*i.e* the quarks and leptons) must be partnered by spin 0 particles (called "sfermions"), not spin 1 bosons. This is because the only spin 1 bosons allowed in a renormalizable theory are the gauge bosons and they have to belong to the adjoin representation of the gauge group, but the quarks and leptons do not. Instead, the gauge bosons are partnered by new spin- $\frac{1}{2}$ "gauginos" (spin- $\frac{3}{2}$ being rules out by the requirement of renormalizability). The need to introduce new supersymmetry partners, rather than interrelate the known bosons and fermions, is undoubtedly a major setback for supersymmetry.

For completeness, we briefly consider also supermultiplets of particles with nonzero mass M . in the particle's rest frame, $P_\mu = (M, 0, 0, 0)$ so the anticommutator (2.27) becomes

$$\{\psi_a, \psi_b^+\} = 2M\delta_{ab}. \quad (2.32)$$

We see that $\frac{\psi_a^+}{\sqrt{2M}}$ and $\frac{\psi_a}{\sqrt{2M}}$ act as creation and annihilation operators, repetitively, for both $a = 1$ and 2 . Starting from a spin state $|s, s_3\rangle$, which is annihilated by the ψ_a , we reach three other states by the action of ψ_1^+, ψ_2^+ , and $\psi_1^+\psi_2^+ = -\psi_2^+\psi_1^+$.

2.5 The Wess-Zumino Model

We are now going to consider the construction of supersymmetric field theories. We shall begin with the model of Wess and Zumino (1974) of the massless spin 0, spin- $\frac{1}{2}$ multiplet. Indeed, probably the most intuitive way of introducing supersymmetry is to explore, through this simple model, possible *Fermi-Bose* symmetries of the Lagrangian. It could therefore equally well have been the starting point for our discussion of supersymmetry.

The simplest multiplet in which to search for supersymmetry consists of a two-component *Weyl spinor* (or equivalently a four-component of *Majorana spinor*) together with two real scalar fields. To be specific, we take a massless Majorana spinor field ψ , and massless scalar and pseudoscalar fields A and B , respectively. The Kinetic energy is

$$\mathcal{L} = \frac{1}{2}(\partial^\mu A)(\partial_\mu A) + \frac{1}{2}(\partial^\mu B)(\partial_\mu B) + \frac{1}{2}\bar{\psi}\gamma_\mu\partial^\mu\psi. \quad (2.33)$$

The unfamiliar factor of $\frac{1}{2}$ in the fermion term arises because ψ is a Majorana spinor; a Dirac spinor is a linear combination of two Majorana spinors (see (2.22)).

The following bilinear identities are particularly useful when exploring supersymmetry. For any two Majorana spinors ψ_1, ψ_2 we have

$$\bar{\psi}_1 \Gamma \psi_2 = \eta \bar{\psi}_2 \Gamma \psi_1 \quad (2.34)$$

where $\eta = (1, 1, -1, 1, -1)$ for $\Gamma = (1, \gamma_5, \gamma_\mu, \gamma_\mu \gamma_5, \sigma_{\mu\nu})$.

To discover the Fermi-Bose symmetries of $\mathfrak{L}$, we make the following infinitesimal transformations: $A \rightarrow A' = A + \delta A$, etc., where

$$\delta A = \bar{\varepsilon} \psi, \quad (2.35a)$$

$$\delta B = i \bar{\varepsilon} \gamma_5 \psi, \quad (2.35b)$$

$$\delta \psi = -i \gamma^\mu \partial_\mu (A + i \gamma_5 B) \varepsilon. \quad (2.35c)$$

ε being a constant infinitesimal Majorana spinor that anticommutes with ψ and commutes with A and B . These transformations are clearly Lorentz-covariant but otherwise δA and δB are just fairly obvious first guesses. The possibility of just constructing two independent invariant quantities $\bar{\varepsilon} \psi$ and $\bar{\varepsilon} \gamma_5 \psi$ is the reason for introducing both scalar and pseudoscalar fields. Since A and ψ have mass dimensions 1 and $\frac{3}{2}$, respectively, ε must have dimension $-\frac{1}{2}$. The derivative in $\delta \psi$ is therefore required to match these dimensions. We have assumed that the transformations have to be linear in the fields.

Under (2.35) the change in $\mathfrak{L}$ can be written in the form

$$\begin{aligned}
\delta\mathfrak{L} &= \partial^\mu A \partial_\mu(\delta A) + \partial^\mu B \partial_\mu(\delta B) + \frac{1}{2}i(\delta\bar{\psi})\gamma^\mu\partial_\mu\psi + \frac{1}{2}i\bar{\psi}\gamma^\mu\partial_\mu(\delta\psi) \\
&= \partial^\mu A \bar{\varepsilon}\partial_\mu\psi + i\partial^\mu B \bar{\varepsilon}\gamma_5\partial_\mu\psi - \frac{1}{2}\bar{\varepsilon}\gamma^\nu\gamma^\mu\partial_\mu(A + i\gamma_5 B)\partial_\nu\psi + \frac{1}{2}\bar{\psi}\gamma^\mu\partial_\mu[\gamma^\nu\partial_\nu(A + i\gamma_5 B)\varepsilon] \\
&= \partial_\mu \left[\bar{\varepsilon}\{\partial^\mu(A + i\gamma_5 B) - \frac{1}{2}\gamma^\nu\gamma^\mu\partial_\nu(A + i\gamma_5 B)\}\psi \right] \\
&= \partial_\mu \left[\frac{1}{2}\bar{\varepsilon}\gamma^\mu\{\partial(A + i\gamma_5 B)\}\psi \right], \tag{2.36}
\end{aligned}$$

where we have used the identities

$$\bar{\varepsilon}\psi = \psi\bar{\varepsilon}, \text{ and } \bar{\varepsilon}\gamma^5\psi = \bar{\psi}\gamma^5\varepsilon, \tag{2.37}$$

of (2.34). Since $\delta\mathfrak{L}$ is a total derivative, it integrates to zero when we form the action. hence, the action is invariant under the combined global supersymmetric transformations (2.35) that mix the fermion and the boson fields. As usual, "global" is used to indicate that ε is independent of space-time.

We have remarked that the $\delta\psi$ transformations (2.35c) contains a derivative. It thus relates the Fermi-Bose symmetry to the Poincare' group. In particular, the appearance of the time derivative gives an *absolute* significance to the total energy, which is normally absent in theories that do not involve gravity.

Returning attention again to the global transformations (2.35), we recall that the commutator of two successive transformations of a symmetry group must itself be a symmetry transformation. In this way we identify the algebra of the generators of the group. To obtain the corresponding result for supersymmetry, we must therefore

consider two successive supersymmetric transformations like (2.35). For example, if for the scalar field A we may take a transformation (2.35a) associated with parameter ε_1 , followed another with parameter ε_2 , then we obtain from (2.35c)

$$\delta_2(\delta_1 A) = \delta_2(\bar{\varepsilon}_1 \psi) = -i\bar{\varepsilon}_1 \gamma^\mu \partial_\mu (A + i\gamma_5 B) \varepsilon_2. \quad (2.38)$$

Hence, the commutator

$$\begin{aligned} (\partial_2 \partial_1 - \partial_1 \partial_2) A &= -i\bar{\varepsilon}_1 \gamma^\mu \partial_\mu (A + i\gamma_5 B) \varepsilon_2 + i\bar{\varepsilon}_2 \gamma^\mu \partial_\mu (A + i\gamma_5 B) \varepsilon_1 \\ &= -2i\bar{\varepsilon}_1 \gamma^\mu \varepsilon_2 \partial_\mu A \\ &= -2\bar{\varepsilon}_1 \gamma^\mu \varepsilon_2 P_\mu A \end{aligned} \quad (2.39)$$

as the terms involving B cancel when we use the identities for Majorana spinors (2.34) and $i\partial_\mu = P_\mu$.

Now the generator of supersymmetric transformations Q_α is a four-component Majorana spinor, which we define by the requirement that

$$\delta A = \bar{\varepsilon} Q A, \quad (2.40)$$

To make this consistent with (2.39), we form the commutator

$$\begin{aligned}
[\partial_2, \partial_1] A &= [\bar{\varepsilon}_2 Q, \bar{\varepsilon}_1 Q] A = [\bar{Q} \varepsilon_2, \bar{\varepsilon}_1 Q] A \\
&= (\bar{Q}_\beta \varepsilon_{2\beta} \bar{\varepsilon}_{1\alpha} Q_\alpha - \bar{\varepsilon}_{1\alpha} Q_\alpha \bar{Q}_\beta \varepsilon_{2\beta}) A \\
&= -\bar{\varepsilon}_{1\alpha} \varepsilon_{2\beta} \{Q_\alpha, \bar{Q}_\beta\} A,
\end{aligned} \tag{2.41}$$

using (2.37). Writing (2.39) in component form and equating it with (2.41) reveals the requirement

$$\{Q_\alpha, \bar{Q}_\beta\} = 2(\gamma^\mu)_{\alpha\beta} P_\mu \tag{2.42}$$

which is indeed part of the supersymmetry algebra (2.27). The same commutator is found on applying successive supersymmetry transformations to field B .

Finally, we must check that the algebra closes when acting on the spinor field ψ . We find from (2.35c)

$$\begin{aligned}
[\delta_2, \delta_1] \psi &= -i\gamma^\mu \partial_\mu \delta_2 (A + i\gamma_5 B) \varepsilon_1 - (1 \leftrightarrow 2) \\
&= -i\bar{\partial}(\bar{\varepsilon}_2 \psi \varepsilon_1 + i\gamma_5 \bar{\varepsilon}_2 i\gamma_5 \psi \varepsilon_1) \varepsilon_1 - (1 \leftrightarrow 2) \\
&= -2i\bar{\varepsilon}_1 \gamma^\mu \varepsilon_2 \partial_\mu \psi + i\bar{\varepsilon}_1 \gamma^\nu \varepsilon_2 \gamma_\nu \bar{\partial} \psi,
\end{aligned} \tag{2.43}$$

where the last equality uses a Fierz rearrangements to bring the two ε together, as well as the Majorana identities (2.37). If we use the field equation $\bar{\partial}\psi = 0$ for a free massless fermion, the last term vanishes identically and (2.43) has exactly the same form as (2.39) for the field A , and hence we obtain (2.42) as before.

However, there is a problem with (2.43) because it gives the required closure only when ψ satisfies the Dirac equation, but not for interacting fermions that are “off-the-mass-shell”. The reason is that for off-mass-shell particles, the numbers of fermion and boson degrees of freedom no longer match up. A and B still have two bosonic degrees of freedom, whereas the Majorana spinor ψ has four. We can restore the symmetry by adding two extra bosonic fields, $\mathfrak{F}$ and $\mathfrak{G}$ (called auxiliary fields), whose free Lagrangian takes the form

$$\mathfrak{L} = \frac{1}{2}\mathfrak{F}^2 + \frac{1}{2}\mathfrak{G}^2. \quad (2.44)$$

This gives the field equations $\mathfrak{F} = \mathfrak{G} = 0$, so these new fields have no on-mass-shell states. From (2.44) they clearly must have mass dimension 2, so on dimensional grounds their supersymmetry transformations can only take the forms

$$\delta\mathfrak{F} = -i\bar{\varepsilon}\gamma^\mu\partial_\mu\psi, \quad \delta\mathfrak{G} = \bar{\varepsilon}\gamma^5\gamma^\mu\partial_\mu\psi \quad (2.45a)$$

and (2.35c) becomes,

$$\delta\psi = -i\gamma^\mu\partial_\mu(A + i\gamma_5 B)\varepsilon + (\mathfrak{F} + i\gamma_5\mathfrak{G})\varepsilon. \quad (2.45b)$$

The mass dimensions prevent $\mathfrak{F}$ and $\mathfrak{G}$ from occurring in

$$\delta A = \bar{\varepsilon}\psi, \quad \delta B = i\bar{\varepsilon}\gamma_5\psi. \quad (2.45c)$$

Under these modified supersymmetry transformations (2.45), we can show that the unwanted term in (2.43) cancels and, moreover, that

$$[\delta_1, \delta_2] \mathfrak{F} = -2i\bar{\varepsilon}_1 \gamma^\mu \bar{\varepsilon}_2 \partial_\mu \mathfrak{F}, \quad (2.46)$$

and similarly for $\mathfrak{G}$, as required by (2.42).

In this way, we have obtained the spin 0, spin- $\frac{1}{2}$ realization of supersymmetry originally found by Wess and Zumino (1974).

2.6 Mass *and* Interaction Terms *in the* Lagrangian

We have found that the free Lagrangian

$$\mathfrak{L} = \frac{1}{2} \partial_\mu A \partial_\mu A + \frac{1}{2} \partial^\mu B \partial_\mu B + \frac{1}{2} i \bar{\psi} \not{\partial} \psi + \frac{1}{2} \mathfrak{F}^2 + \frac{1}{2} \mathfrak{G}^2, \quad (2.47)$$

that describes the multiplet $(A, B, \psi, \mathfrak{F}, \mathfrak{G})$, is invariant (up to a total derivative) under the supersymmetry transformations (2.45). However, it is easy to check that supersymmetry invariance is still preserved if the Lagrangian is extended to include a quadratic mass term of the form

$$\mathfrak{L}_m = m(\mathfrak{F}A + \mathfrak{G}B - \frac{1}{2} \psi \bar{\psi}) \quad (2.48)$$

and a cubic interaction term

$$\mathfrak{L}_i = \frac{g}{\sqrt{2}} [\mathfrak{F}A^2 - \mathfrak{F}B^2 + 2\mathfrak{G}AB - \bar{\psi}(A - i\gamma_5 B)\psi] \quad (2.49)$$

Higher-order terms must be excluded because they are nonrenormalizable. When we use the classical equations of motion,

$$\frac{\partial \mathfrak{L}}{\partial \mathfrak{F}} = \frac{\partial \mathfrak{L}}{\partial \mathfrak{G}} = 0, \quad (2.50)$$

for the complete Lagrangian, $\mathfrak{L} = \mathfrak{L}_0 + \mathfrak{L}_m + \mathfrak{L}_i$, we find

$$\mathfrak{F} + mA + \frac{g}{\sqrt{2}}(A^2 - B^2) = 0, \quad (2.51a)$$

$$\mathfrak{G} + mB + \sqrt{2}gAB = 0. \quad (2.51b)$$

These equations of motion are purely algebraic and so the dynamics is unchanged if we use them to eliminate the auxiliary fields $\mathfrak{F}$ and $\mathfrak{G}$ from the Lagrangian. We obtain

$$\begin{aligned} \mathfrak{L} = & \frac{1}{2}\partial_\mu A \partial^\mu A + \frac{1}{2}\partial_\mu B \partial^\mu B + \frac{1}{2}\bar{\psi}\mathfrak{F}\psi - \frac{1}{2}m\bar{\psi}\psi - \frac{1}{2}m^2(A^2 + B^2) - \frac{1}{\sqrt{2}}mgA(A^2 + B^2) \\ & - \frac{1}{2}g^2(A^2 + B^2)^2 - \frac{1}{\sqrt{2}}g\bar{\psi}(A - i\gamma^5 B)\psi \end{aligned} \quad (2.52)$$

Several features of this Lagrangian, which are characteristics of supersymmetric theories, are worth nothing. The masses of the scalars and the fermions are all equal. There are cubic and quartic couplings between the scalar fields, and also a Yukawa-type interaction between the fermion ψ and the scalars A , and B yet in total there are only two free parameters: m and g . These interrelation between fermion and boson masses and couplings is the essence of the supersymmetry.

The model can also be shown to have some remarkable renormalization properties in that, despite the presence of the scalar fields, there is no renormalization of the mass and coupling constant (although wave function renormalization is still necessary). the divergences arising from boson loops are cancelled by those from fermion loops which have the opposite sign. This is just the type of cancellation we need to stabilize the gauge hierarchy. These powerful nonrenormalization theorems make supersymmetry particularly compelling. However when we break supersymmetry, as well give the absence of fermion-boson mass degeneracy in nature, we have to be careful to preserve the relations between the couplings of particles of different spin implied in (2.52).

2.7 The Superpotential

To see how this results generalize with higher symmetries, it is convenient to work entirely with left handed fermion fields. Recall from (2.29) that Majorana spinor ψ can be formed entirely from a left-handed Weyl spinor

$$\psi = \psi_L + C\bar{\psi}_L^T \quad (2.53)$$

and that the mass term is

$$\begin{aligned} \mathcal{L} &= m\bar{\psi}\psi = m\bar{\psi}_R^c\psi_L + \text{herm.conj.} \\ &= m\psi_L^T C\psi_L + \text{herm.conj.} \end{aligned} \quad (2.54)$$

using the relation

$$\overline{\psi}_R^c = \psi_R^{c+} \gamma^0 = \psi_L^{*+} \gamma_0^{T+} C^+ \gamma^0 = -\psi_L^T C^{-1} = \psi_L^T C. \quad (2.55)$$

For simplicity we have set $-C^{-1} = C$, which is valid in all the familiar representations of the Dirac matrices. We can rewrite the supersymmetry Lagrangian of section (2.6) using just a left handed field ψ_L , and complex fields ϕ and F for its scalar partner, viz.,

$$\phi \equiv \frac{1}{\sqrt{2}}(A + iB), \quad \text{and} \quad F \equiv \frac{1}{\sqrt{2}}(\mathfrak{F} - i\mathfrak{G}) \quad (2.56)$$

From $\mathfrak{L} = \mathfrak{L}_0 + \mathfrak{L}_m + \mathfrak{L}_i$ we obtain

$$\mathfrak{L} = \partial_\mu \phi^* \partial^\mu \phi + i \overline{\psi}_L \mathfrak{D} \psi_L + F F^* + m(\phi F - \frac{1}{2} \psi_L^T C \psi_L) + \text{herm.conj.} + g(\phi^2 F - \phi \psi_L^T C \psi_L) + \text{hem.conj.} \quad (2.57)$$

Then using the equation of motion $\frac{\partial \mathfrak{L}}{\partial \mathfrak{F}} = 0$, which gives

$$F^* = -m\phi - g\phi^2, \quad (2.58)$$

we can eliminate the auxiliary field F^* and so the Lagrangian becomes

$$\mathfrak{L} = \partial_\mu \phi^* \partial^\mu \phi + i \overline{\psi}_L \mathfrak{D} \psi_L - |m\phi + g\phi^2| - (\frac{1}{2} m \psi_L^T C \psi_L + g \phi \psi_L^T C \psi_L + \text{herm.conj.}) \quad (2.59)$$

It is useful to re-examine the Lagrangian (2.57) in terms of an analytic function $W(\phi)$, known as the “superpotential”, viz.,

$$\mathfrak{L} = \mathfrak{L}_{K.E.} + FF^* + F^* \frac{\partial W}{\partial \phi} + F \frac{\partial W^*}{\partial \phi^*} - \frac{1}{2} \left(\frac{\partial^2 W}{\partial \phi^2} \psi_L^T C \psi_L + h.c. \right), \quad (2.60)$$

where $\mathfrak{L}_{K.E.}$ denotes the sum of the kinetic energy terms of the ϕ and ψ_L fields.

Note that W , which is of dimension 3, depends only on ϕ and not on ϕ^* . Upon using

$\frac{\partial \mathfrak{L}}{\partial \mathfrak{F}} = \frac{\partial \mathfrak{L}}{\partial \mathfrak{F}^*} = 0$ to eliminate the auxiliary fields, we find

$$\mathfrak{L} = \mathfrak{L}_{K.E.} - \left| \frac{\partial W}{\partial \phi} \right|^2 - \frac{1}{2} \left(\frac{\partial^2 W}{\partial \phi^2} \psi_L^T C \psi_L + h.c. \right). \quad (2.61)$$

For a renormalizable theory W can be, at most, a cubic function of ϕ , since otherwise the Lagrangian would contain couplings with dimension less than 0. Substituting

$$W = \frac{1}{2} m \phi^2 + \frac{1}{3} g \phi^3 \quad (2.62)$$

into (2.61) immediately reproduces the Lagrangian of (2.59). The superpotential is the only free function in the supersymmetry Lagrangian and determines both the potential of the scalar fields, and the masses and couplings of fermions and bosons.

In general there may be several chiral multiplets to consider, For example, if ψ^i belongs to a representation of an $SU(N)$ symmetry group, we will have the supermultiplets

$$(\phi^i, \psi_L^i) \quad (2.63)$$

where in the fundamental representation $i = 1, 2, \dots, N$. From (2.61) we readily obtain a Lagrangian that is invariant under the additional symmetry and incorporates the new

supermultiplets. It is

$$\mathcal{L}_{chiral} = \sum_i |\partial_\mu \phi^i|^2 + i \sum_i \overline{\psi_L^i} \not{\partial} \psi_L^i - \sum_i \left| \frac{\partial W}{\partial \phi^i} \right|^2 - \frac{1}{2} \left(\sum_{i,j} \frac{\partial^2 W}{\partial \phi^i \partial \phi^j} \psi_L^{iT} C \psi_L^j + \text{herm.conj.} \right), \quad (2.64)$$

and the most general form of the superpotential W is

$$W = \lambda_i \phi^i + \frac{1}{2} m_{ij} \phi^i \phi^j + \frac{1}{3} g_{ijk} \phi^i \phi^j \phi^k, \quad (2.65)$$

where the coefficients m and g are completely symmetric under interchange of indices. The relevance of the term that is linear in the fields will be discussed below. Since W must be invariant under $SU(N)$ symmetry transformations this term can only occur if a gauge-singlet field exists.

2.8 Supersymmetric Gauge Theory

A combination of supersymmetry with gauge theory is clearly necessary if these ideas are to make any contact with the real world. In addition to the chiral multiplets of (2.63), we must include the “gauge” supermultiplets

$$(A_\mu^a, \chi^a), \quad a = 1, 2, 3, \dots, N^2 - 1, \quad (2.66)$$

where A_μ^a are the spin 1 gauge bosons of the gauge group G (taken to be $SU(N)$) and χ^a are their Majorana fermion superpartners (the so called “gauginos”). These boson-fermion pairs, which in the absence of symmetry breaking are assumed to be

massless, belong to the adjoint representation of the gauge group. Our task is to find a supersymmetry, and a gauge-invariant Lagrangian containing all these chiral and gauge supermultiplets.

The gauge multiplets are described by the Lagrangian

$$\mathfrak{L}_G = \frac{-1}{4} F_{\mu\nu}^a F_a^{\mu\nu} + \frac{1}{2} i \overline{\chi^a} (D\chi)_a + \frac{1}{2} (D^a)^2, \quad (2.67)$$

where the gauge field-strength tensor is

$$F_a^{\mu\nu} = \partial^\mu \chi_a - g_G f_{abc} A_b^\mu A_c^\nu, \quad (2.68)$$

D^μ is the covariant derivative satisfying

$$(D^\mu \chi)_a = \partial^\mu \chi_a - g_G f_{abc} A_b^\mu \chi_c \quad (2.69)$$

and D^μ are auxiliary scalar fields (similar to F_i of the chiral multiplet).

Actually, for this pure gauge Lagrangian the equation of motion $\partial \mathfrak{L}_G / \partial D^a = 0$, implies $D^a = 0$; however, it will become nonzero when the chiral fields are coupled in the notation will be familiar: g_G and f_{abc} are the coupling and the structure constants of the gauge group, and in (2.96) the matrices T^b representing the generators in the adjoint representation have been replaced by $(T^b) = if_{abc}$. It is straight forward to show that the lagrangian $\mathfrak{L}_G$ is invariant, and that the algebra closes, under the supersymmetry transformation

$$\begin{aligned}
\delta A_a^\mu &= -\bar{\varepsilon}\gamma^\mu\gamma^5\chi_a, \\
\delta\chi^a &= -\frac{1}{2}\sigma^{\mu\nu}F_{\mu\nu}^a\gamma^5\varepsilon + D^a\varepsilon, \\
\delta D^a &= -i\bar{\varepsilon}(D\chi)^a,
\end{aligned} \tag{2.70}$$

where ε is a constant infinitesimal Majorana spinor. This transformation is analogous to (2.35) for the chiral multiplet.

To include the chiral fields (ϕ^i, ψ_L^i) , we add $\mathfrak{L}_{chiral}$ of (2.64) but substituting the covariant derivative D_μ for ∂_μ in the kinetic energy terms, viz.,

$$\partial_\mu \rightarrow D_\mu = \partial_\mu + ig_G T^a A_\mu^a, \tag{2.71}$$

where T^a are the matrices representing the generators of the gauge group in the representation to which (ϕ_i, ψ_i) belong. To ensure the supersymmetry of the combined “chiral + gauge” Lagrangian, we must include two further terms, and write

$$\mathfrak{L} = \mathfrak{L}_{chiral} + \mathfrak{L}_G - g_G \phi_i^* (T^a)_{ij} \phi_j D^a + [\sqrt{2} g_G \phi_i^* \bar{\chi}^a (T^a)_{ij} P_L \psi_j + herm.conj.] \tag{2.72}$$

where $P_L \equiv \frac{1}{2}(1 - \gamma^5)$, and replace ∂_μ in (2.45) by D_μ . Using $\partial\mathfrak{L}_G/\partial D^a = 0$ to eliminate the auxiliary field gives

$$D^a = g_G \phi_i^* (T^a)_{ij} \phi_j. \tag{2.73}$$

The term in the Lagrangian that contribute to the potential for the scalar fields are evident by inspection of (2.61) and (2.67). They are

$$\begin{aligned} V(\phi, \phi^*) &= |F_i|^2 + \frac{1}{2} D_a^2 \\ &= \sum_i \left| \frac{\partial W}{\partial \phi_i} \right|^2 + \frac{1}{2} \sum_a [g_G \sum_{i,j} \phi_i^* (T^a)_{ij} \phi_j]^2, \end{aligned} \quad (2.74)$$

which are known as F and D terms, respectively. This potential will play a central role in the spontaneous breaking of supersymmetry and the gauge symmetry.

Model building begins with the supersymmetry Lagrangian (2.72). Apart from the choice of gauge group and the representations formed by the chiral multiplets, the only freedom lies in the choice of the superpotential $W(\phi_i)$ which must, of course, be a single of the gauge group.

2.9 Spontaneous Breaking of Supersymmetry

The particles observed in nature show no sign whatsoever of degeneracy between fermions and bosons. Even the photon and neutrino, which appear to be degenerate in mass, can not be supersymmetric partners. Hence, supersymmetry, if it is to be relevant in nature, must be broken.

The breaking could be either explicit or spontaneous. Explicit breaking would be quite *ad hoc*. The supersymmetry generators would no longer commute with the Hamiltonian,

$$[Q_\alpha, H] \neq 0 \tag{2.75}$$

and so the violation would have to be small enough to preserve the good features of supersymmetry and yet large enough to push the supersymmetric partners out of reach of current experiments. However, we would inevitably lose the nice nonrenormalization theorems and, even worse, any attempt to embrace gravity viz local supersymmetry would be prohibited. So, instead, we prefer to consider the spontaneous breaking of supersymmetry, not least because this has proved so successful previously for breaking gauge symmetries. Hence, we assume that the Lagrangian is supersymmetric but that the vacuum state is not, that is

$$[Q_\alpha, H] = 0, \quad \text{but} \quad Q_\alpha |0\rangle \neq 0 \tag{2.76}$$

A new feature arises here, however. The Higgs mechanism of spontaneous symmetric breaking is not available in supersymmetry because, if we were to introduce a spin 0 field with negative mass-squared, its fermionic superpartner would have an imaginary mass. Also, using the anticommutator (2.27),

$$\{Q_\alpha, Q_\delta^+\} \gamma_{\delta\beta}^0 = 2\gamma_{\alpha\beta}^0 P_\mu, \tag{2.77}$$

we can directly establish a general and important theorem. If we multiply (2.77) by $\gamma_{\beta\alpha}^0$ and sum over β and α , we obtain

$$\sum_{\alpha} \{Q_{\alpha}, Q_{\alpha}^{+}\} = 8P_0 = 8H \quad (2.78)$$

and hence

$$8 \langle 0 | H | 0 \rangle = \sum_{\alpha} |Q_{\alpha} | 0 \rangle|^2 + \sum_{\alpha} |Q_{\beta}^{+} | 0 \rangle|^2 \quad (2.79)$$

It follows immediately that

- 1- The vacuum energy must be greater than or equal to zero;
- 2- If the vacuum is supersymmetric, that is, if $Q_{\alpha} | 0 \rangle = Q_{\beta}^{+} | 0 \rangle = 0$ for all α , the vacuum energy is zero; and
- 3- Conversely, if supersymmetry is spontaneously broken, $Q_{\alpha} | 0 \rangle \neq 0$, then the vacuum energy is positive.

These results have a disappointing consequences. Conclusion (1) gives an absolute meaning to the zero of energy, a fact that it was hoped to use to explain why the vacuum energy of the universe is zero or very close to zero. But now from (3) we see that broken supersymmetry implies a positive vacuum energy.

Leaving this aside we can see from (3) that supersymmetry breaking is rather special because it requires the ground-state energy to be positive.

In The classical approximation, the energy of the ground state is given by the minimum of the potential (2.74):

$$\begin{aligned}
V(\phi, \phi^*) &= |F_i|^2 + \frac{1}{2} D_a^2 \\
&= \sum_i \left| \frac{\partial W}{\partial \phi_i} \right|^2 + \frac{1}{2} \sum_{\beta, a} [g_\beta \phi_i^* (T_\beta^*)_{ij} \phi_j + \eta \delta_{\beta 1}]^2, \tag{2.80}
\end{aligned}$$

with

$$W = \lambda_i \phi_i + \frac{1}{2} m_{ij} \phi_i \phi_j + g_{ijk} \phi_i \phi_j \phi_k \tag{2.81}$$

The sum over β has been included to allow for the possibility of different gauge groups with different couplings, and the constant term η can only occur if β labels a $U(1)$ factor.

It is evidently hard to break supersymmetry. The minimum $V = 0$ will occur when $\phi_i = 0$ for all i (and so supersymmetry will be broken) unless one of the following conditions applies.

1- $\lambda_i \neq 0$, that is, there exists a gauge singlet field ϕ_i , so W can contain a linear term yet still be gauge invariant ($F - type$ breaking);

2- $\eta \neq 0$, so the gauge group contain an abelian $U(1)$ factor ($D - type$ breaking). This is a necessary but not a sufficient requirement. This mechanism can not occur in GUT s because they are based on simple gauge groups that do not have $U(1)$ factor.

There is an alternative way of seeing that the spontaneous symmetry breaking of supersymmetry can only be accomplished by $\langle F \rangle \neq 0$ and/or $\langle D \rangle \neq 0$. If we look back at the general structure of supersymmetric transformations (2.45) for the chiral multiplet (ϕ, ψ, F) , which takes the form

$$\delta\phi \sim \psi, \quad \delta\psi \sim \mathfrak{D}\phi + F, \quad \delta F \sim \mathfrak{D}\phi \quad (2.82)$$

and at (2.70) for the gauge multiplet (A_μ, χ, D) , in which

$$\delta A_\mu \sim \gamma_\mu \chi, \quad \delta\chi \sim \sigma^{\mu\nu} F_{\mu\nu} + D, \quad \delta D \sim \mathfrak{D}\chi, \quad (2.83)$$

and note that the vacuum expectation values of the spinor and tensor fields and $\partial_\mu\phi$ must be zero to preserve the Lorentz invariance of the vacuum, then it is only to break the symmetry through the non-zero VEVs of the auxiliary fields F and D .

The spontaneous breaking of supersymmetry requires

$$Q_\alpha |0\rangle \neq 0 \quad (2.84)$$

and $Q_\alpha |0\rangle$ is necessarily a fermionic state, which we denote by $|\psi_G\rangle$.

Since the Q_α commute with H , the state $|\psi_G\rangle$ must be degenerate with the vacuum. It must therefore describe a massless fermion (with zero momentum). The situation is thus exactly analogous to the spontaneous breaking of ordinary global symmetry in which massless Goldstone bosons are created out of the vacuum. Here the spontaneous breaking of global supersymmetry implies the existence of a massless fermion, which is called “Goldstino.”

We next consider examples of these two types of symmetry breaking *F – type* and *D – type* introduced in (1) and (2) above.

2.10 F-type Breaking (O'raifeartaigh Model)

A simple example of supersymmetry breaking arising from the presence of a linear term in the superpotential W is provided by

$$W = -\lambda A + mBC + gAB^2, \quad (2.85)$$

which contain three complex scalar fields A, B , and C . In this example the scalar potential (2.80) becomes

$$\begin{aligned} V &= \left| \frac{\partial W}{\partial A} \right|^2 + \left| \frac{\partial W}{\partial B} \right|^2 + \left| \frac{\partial W}{\partial C} \right|^2 \\ &= |-\lambda + gB^2|^2 + |mC + 2gAB^2|^2 + |mB|^2 \end{aligned} \quad (2.86)$$

and we see that $V = 0$ is excluded because the last term is only zero if $B = 0$, but then the first term is positive-definite. We conclude that $V > 0$ and that supersymmetry is broken. Provided that $m^2 > 2\lambda g$ the potential V has a minimum when $B = C = 0$, independently of the value of A . For simplicity, we set $A = 0$ at the minimum.

As usual, the scalar masses are determined by evaluating

$$V_{AB} \equiv \frac{\partial^2 V}{\partial A \partial B}, \quad \text{etc.}, \quad (2.87)$$

at the minimum. The only non-zero elements are

$$\langle V_{BB'} \rangle B^2 + 2 \langle V_{BB^*} \rangle B B^* + \langle V_{B^* B^*} \rangle B^{*2} = (m^2 - 2g\lambda) B_1^2 + (m^2 + 2g\lambda) B_2^2, \quad (2.88)$$

where $B = (B_1 + iB_2)/\sqrt{2}$, and so the real scalar fields B_1 and B_2 have $(mass)^2 = m^2 \mp 2g\lambda$, respectively.

The fermion masses are obtained by evaluating $\partial^2 W / \partial A \partial B$, etc., at the minimum (see (2.64)). From (2.85) we find that the only non-zero term is

$$\frac{\partial^2 W}{\partial B \partial C} = m \quad (2.89)$$

and so the fermion mass matrix takes the form

$$M_F = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & m \\ 0 & m & 0 \end{pmatrix} \quad (2.90)$$

in the bases of Majorana spinors ψ_A, ψ_B, ψ_C . The massless Goldstino state ψ_A is evident, and the off-diagonal structure signals that the two Majorana spinors ψ_B, ψ_C will combine to give a single Dirac fermion of mass m . Despite the supersymmetry breaking, there is still an equality between the sum of the $(mass)^2$ of the bosons and that of the fermions. Explicitly, for each degree of freedom we have the masses given in table (2.1)

<i>Bosons</i>			<i>Fermions</i>	
A	B	C	ψ_A	ψ_B, ψ_C
$0, 0$	$m^2 \pm 2\lambda g$	m^2, m^2	$0, 0$	m^2, m^2, m^2, m^2

Table (2.1): Masses of bosons and their fermionic partners

Only B suffers supersymmetry breaking. The reason is that it is the only field that couples to the Goldstino; its coupling $gB\bar{\psi}_B\psi_A$ appears when (2.85) is inserted into (2.64). The value of the potential at the minimum can be written

$$\langle V \rangle = \lambda^2 \equiv M_S^4 \quad (2.91)$$

where the mass splitting within the supermultiplet of table (2.1) are therefore

$$\Delta m^2 \approx gM_S^2, \quad (2.92)$$

where g is the coupling to the Goldstino.

This simple model illustrates several more general results. The mass relation is a particular example of the “super-trace relation”,

$$STr(M^2) \equiv \sum_J (2J+1)(-1)^{2J} Tr(M_J^2) = 0, \quad (2.93)$$

which holds whether supersymmetry is spontaneously broken or not. Here M_J is the mass matrix for the fields of spin- J , and the sum is over all the physical particles of the theory. Relation (2.93) holds in lowest-order perturbation theory. We say that it is a “*tree-level*” result because it neglects corrections due to the diagrams containing loops. This super-trace mass relation is important because it ensures that the scalars are not subject to quadratically divergent renormalization.

We may readily verify that (2.93) holds for an arbitrary multiplet structure. If there are several chiral multiplets (ϕ_i, ψ_i) , then it is convenient to arrange the scalar fields

and their complex conjugates as a column vector so that the boson mass terms have the matrix structure

$$\begin{pmatrix} \phi^* & \phi \end{pmatrix} \begin{pmatrix} X & Y \\ Y^+ & X \end{pmatrix} \begin{pmatrix} \phi \\ \phi^* \end{pmatrix}. \quad (2.94)$$

The block diagonal parts of the boson $(mass)^2$ matrix, M_B^2 , have elements

$$\begin{aligned} X_{ij} &= \frac{\partial^2 V}{\partial \phi_i \partial \phi_j^*} = \sum_k \left(\frac{\partial^2 W}{\partial \phi_i \partial \phi_k} \right) \left(\frac{\partial^2 W}{\partial \phi_j^* \partial \phi_k^*} \right) \\ &= \sum_k (M_F)_{ik} (M_F^*)_{kj} = (M_F M_F^*)_{ij} \end{aligned} \quad (2.95)$$

where M_F is the fermion mass matrix and so it follows that

$$Tr(M_B^2) = 2Tr(M_F^2) \quad (2.96)$$

at tree level.

We can also show that the fermion mass matrix has a zero eigenvalue and hence identify the Goldstone. At the minimum of the potential,

$$0 = \frac{\partial V}{\partial \phi_i} = \frac{\partial}{\partial \phi_i} \left(\sum_j \left| \frac{\partial W}{\partial \phi_j} \right|^2 \right) = \sum_j \left(\frac{\partial^2 W}{\partial \phi_i \partial \phi_j} \right) \left(\frac{\partial W}{\partial \phi_j} \right)^* = \sum_j (M_F)_{ij} \langle F_j \rangle^*. \quad (2.97)$$

Thus, the mass matrix M_F annihilates the fermion state

$$\psi_G = \sum_j \langle F_j \rangle^* \psi_j \quad (2.98)$$

which is thus identified as the massless Goldstino. In our example, $\psi_G = \psi_A$ since $\langle F_B \rangle = \langle F_c \rangle = 0$.

However, the equality (2.96), which is so desirable to ensure the boson-fermion loop cancellations, is not supported experimentally. The difficulty is that in these simple models, the relation applies to each supermultiplet separately. Hence, for the electron, for example, we require

$$2m_e^2 = m_A^2 + m_B^2, \quad (2.99)$$

which implies that one of the two scalar electrons (A, B) must have a mass less than or equal to that of the electron. Such a particle would have been detected long time ago if it existed.

2.11 D-type Breaking (The Fayet-Iliopoulos Model)

As a simple example of supersymmetry breaking caused by the presence of a $U(1)$ factor in the gauge group, we take a supersymmetric version of QED with two chiral multiplets (ϕ_+, ψ_+) and (ϕ_-, ψ_-) , where the subscripts give the sign of the charge. The $U(1)$ gauge-invariant superpotential is

$$W = m\phi_+\phi_- \quad (2.100)$$

and so the scalar potential (2.80) becomes

$$\begin{aligned}
V &= m^2 |\phi_+|^2 + m^2 |\phi_-|^2 + \frac{1}{2} \left[e \left(|\phi_+|^2 - |\phi_-|^2 \right) + \eta \right]^2 \\
&= \frac{1}{2} e^2 \left(|\phi_+|^2 - |\phi_-|^2 \right)^2 + (m^2 + e\eta) |\phi_+|^2 + (m^2 - e\eta) |\phi_-|^2 + \frac{1}{2} \eta^2
\end{aligned} \tag{2.101}$$

Various possible forms for V are shown in Figure (2.1)

The Pic

Provided $m^2 > 2\eta$ (where $e\eta > 0$), the minimum occurs at

$$\phi_+ = \phi_- = 0, \tag{2.102}$$

so $U(1)$ gauge invariance is not spontaneously broken, but supersymmetry is broken since $V \neq 0$. The boson masses are split, $m_{\pm}^2 = m^2 \pm e\eta$, whereas the fermion masses are unaffected by the breakdown of supersymmetry. Like (2.90) the (off-diagonal) form of the fermion mass matrix in the ψ_+ , ψ_- Majorana basis implies that these two states combine together to give a Dirac fermion of mass m . The fermion-boson mass splitting signals that the breakdown of supersymmetry but the $(mass)^2$ equality still holds, since

$$m_+^2 + m_-^2 = 2m^2 \tag{2.103}$$

For $m^2 > e\eta$, the $U(1)$ symmetry is unbroken and the gauge multiplet (A_μ, χ) remains massless. The fermion χ is the “Goldstino” arising from the spontaneous supersymmetry breaking.

The case $m^2 < e\eta$ is more interesting. The minimum of the potential now occurs at

$$\phi_+ = 0 \qquad \phi_- = v \qquad (2.104)$$

where $e^2 v^2 = (e\eta - m^2)$. Now, both the $U(1)$ gauge symmetry and supersymmetry are spontaneously broken; see Figure (2.1.c). We find that the complex field ϕ_+ has $(mass)^2 = 2m^2$, while one component of ϕ_- is “eaten” by the usual Higgs mechanism to give $(mass)^2 = 2e^2 v^2$ to the vector gauge field A_μ , and the remaining component also acquires $(mass)^2 = 2e^2 v^2$. A linear combination of the ψ_+ and χ Majorana fields forms the massless “Goldstino”, whereas the two remaining combinations of ψ_+, ψ_- , and χ both have $(mass)^2 = m^2 + 2e^2 v^2$. Despite the symmetry breaking, the super-trace mass relation (2.93) remains true, that is

$$2(2m^2) + 2e^2 v^2 - (2+2)(m^2 + 2e^2 v^2) + 3(2e^2 v^2) \qquad (2.105)$$

It is straight forward to show that, as in sec. (2.9), the super-trace relation $STr(M^2) = 0$ holds in general, with just one exception, and that the Goldstino can be identified as the combination

$$\psi_G = \langle F_j \rangle \psi_j - \frac{1}{\sqrt{2}} \langle D^a \rangle \chi_a. \qquad (2.106)$$

Since the super-trace relation leads to problems, as we found in (2.99), it is desirable to explore the exception. In the presence of a $U(1)$ factor of the gauge group, we find that (2.93) becomes

$$STr (M^2) = 2 \langle D \rangle Tr Q, \quad (2.107)$$

where Q is the $U(1)$ charge matrix of the chiral multiplet and D is the auxiliary field. Perhaps this extra contribution will permit the superpartners to be sufficiently massive to escape detection. Unfortunately, the $U(1)$ of the standard model is not suitable, as the weak hypercharge Y must satisfy $Tr Y = 0$. This extra contribution is also absent in GUT s, which have no $U(1)$ factor. To have an additional $U(1)$ with $Tr Q \neq 0$ would create new problems with triangle anomalies, which can only be avoided by introducing new chiral multiplets. Thus far, no satisfactory model of $D - type$ breaking has been found.

2.12 The Supersymmetric Standard Model

The standard model (SM) has 28 bosonic degrees of freedom (12 massless gauge bosons and 2 complex scalars) together with 90 fermionic degrees of freedom (3 families each with 15 two-component Weyl fermions). To make the model supersymmetric we must clearly introduce additional particles. In fact, since none of the observed particles pare off,

we have to double the number. In section 2.3 we saw that the gauge bosons are partnered by spin- $\frac{1}{2}$ gauginos, and these cannot be identified with any of the quarks and leptons. So the latter have to be partnered by new spin 0 squarks and sleptons.

We also have to complete the Higgs supermultiplet. now the $Y = -1$ Higgs doublet

has the same quantum numbers as the $(\nu, e^-)_L$ doublet, so one might try to identify the Higgs with a spin 0 slepton. Unfortunately even this is not possible, because any attempt to partner a lepton with a Higgs, by giving the latter a non-zero lepton number L , leads to L -violating processes and large $\Delta L = 2$ Majorana mass terms. Even worse, in the standard Higgs (ϕ) generates masses for the down-type quarks and the charged leptons, while its charge conjugate $(\phi_c = i\tau_2\phi^*)$ gives masses to the up- type quarks. Now the superpotential W is a function only of ϕ and not ϕ^* , and so in supersymmetry we need to introduce a second unrelated Higgs doublet. There is an alternative way to see this. Under charge conjugation, the helicity of the spin- $\frac{1}{2}$ partner of the Higgs (“the Higgsino”) is reversed, and so it proves impossible to use a single Higgs to give masses to both up-type and down-type quarks. the second (complex) doublet is also needed to cancel the anomalies that would arise if there were only on Higgsino. As in the standard model, three of the Higgs fields are absorbed to make the $W^\pm$ and Z bosons massive, and we are therefore left with two charged and three neutral massive Higgs particles.

The particle content of the supersymmetric standard model is shown in table (2.2) .

<i>Chiral Multiplets</i>		<i>Gauge Multiplets</i>	
Spin-$\frac{1}{2}$	Spin 0	Spin 1	Spin-$\frac{1}{2}$
Quark q_L, q_R	Squark $\tilde{q}_L, \tilde{q}_R$	photon γ	photino $\tilde{\gamma}$
Lepton l_L, l_R	Slepton $\tilde{l}_L, \tilde{l}_R$	W, Z bosons	Wino $\tilde{W}$, Zino $\tilde{Z}$
Higgsino $\tilde{\phi}, \tilde{\phi}'$	Higgs ϕ, ϕ'	Gluon g	Gluino $\tilde{g}$

Table (2.2) Particle Multiplets in the Supersymmetric Standard Model

There is no doubt that this table is a setback for supersymmetry. To be economical, supersymmetry ought to unite the known fermionic “matter” *quarks* and *leptons* with the vector “forces” γ, g, W, Z , but we have been compelled to keep them separate and to introduce a new superpartner for each particle. A great deal of effort has gone into the search for these superpartners but so far none has been found.

CHAPTER 3

MINIMAL SUPERSYMMETRIC STANDARD MODEL

3.1 Introduction

The minimal supersymmetric extension of the standard model (MSSM) [11] was generated by taking the standard model (SM) and adding the corresponding supersymmetric partners. In addition, the MSSM contains two hypercharge $Y = \pm 1$ Higgs doublets, which is the minimal structure for the Higgs sector of an anomaly-free supersymmetric extension of the standard model. The supersymmetric structure of the theory also requires (at least) two Higgs doublets to generate mass for both up-type and down-type quarks (and charged leptons). All renormalizable supersymmetric interactions consistent with (global) $B - L$ conservation ($B = \text{baryon number}$ and $L = \text{lepton number}$) are included. Finally, the most general soft-supersymmetric-breaking terms are added.

If supersymmetry is relevant for explaining the scale of electroweak interactions, then the mass parameters exist due to the absence of supersymmetric-particle at current accelerators. Additional constraints arise from limits on the contributions of virtual supersymmetric particle exchange to a variety of SM processes.

As a consequence of $B - L$ invariance, the MSSM processes a discrete $R - \text{parity}$

invariance, where $R = (-1)^{3(B-L)-2S}$ for a particle of spin S . This formula implies that all the ordinary SM particles have *even* $-R$ *parity*, whereas the corresponding supersymmetric partners have odd $R - \text{parity}$. The conservation of $R - \text{parity}$ in scattering and decay processes has a crucial impact on supersymmetric phenomenology. For example starting from an initial state involving ordinary ($R - \text{even}$) particles, it follows that supersymmetric particles must be produced in pairs. In general, these particles are highly unstable and decay quickly into lighter states. However R -parity invariance also implies that the lightest supersymmetric particle (LSP) is absolutely stable, and must eventually be produced at the end of a heavy unstable supersymmetric particle.

In order to be consistent with cosmological constraints, the LSP is almost certainly electrically and color neutral. Consequently, the LSP is weakly-interacting in ordinary matter, *i.e.* it behaves like a stable heavy neutrino and will escape detectors without being directly observed. Thus, the canonical signature for ($R - \text{parity}$ conserving) supersymmetric theories is a missing (transverse) energy, due to the escape of the LSP.

Some model builders attempt to relax the assumption of $R - \text{parity}$ conservation. Models of this type must break $B - L$ conservation and are therefore constrained by experiment. Nevertheless, it is still important to allow the possibility of $R - \text{parity}$ violation processes in the search for supersymmetry. In such models the LSP is unstable and supersymmetric particles can be singly produced and destroyed in association with B and L violation. These features lead to a phenomenology of broken $R - \text{parity}$ models that is very different from that of the MSSM

In the MSSM, supersymmetry breaking is accompanied by including the soft-supersymmetry breaking terms. These terms parameterize our ignorance of the fundamental mechanism of supersymmetry breaking. If this breaking occurs spontaneously, then (in the absence) of supergravity a massless goldstone fermion is called the *goldstino* ($\tilde{G}$) must exist. The goldstino would be the LSP and could play an important role in supersymmetric phenomenology. In models that incorporates supergravity (SUGRA), this picture changes. If supergravity is spontaneously broken, the goldstino is absorbed (eaten) by the *gravitino*, the spin- $\frac{3}{2}$ partner of the graviton. By this super-Higgs mechanism, the gravitino acquires a mass. In many models, the gravitino mass is of order as the order of the electroweak-breaking scale, while its coupling are gravitational in strength. Such a gravitino would play no role in supersymmetric phenomenology at colliders. The parameters of the MSSM are conveniently described by considering separately the supersymmetric conserving sector and the supersymmetry breaking sector. Among the parameters of the supersymmetry conserving sector are:

1- gauge couplings: g' , g , and g_s , corresponding to U(1), SU(2), and SU(3) subgroups of the SM respectively;

2- Higgs-Yukawa couplings: λ_e , λ_u , and λ_d (which are 3×3 matrices in flavor space);
and

3- a supersymmetry-conserving Higgs mass parameter μ .

The supersymmetric-breaking sector contains the following set of parameters:

i- gauging Majorana masses M_1 , M_2 and M_3 associated with the U(1), SU(2), and

SU(3) subgroups of the SM;

ii- scalar mass matrices for the squarks and sleptons.;

iii- Higgs-squark trilinear interaction terms (the so-called $A - parameters$) and corresponding terms involving the sleptons; and

iv- three scalar Higgs mass parameters- two-diagonal and one off-diagonal mass terms for two Higgs doublets. These three mass parameters can be re-expressed in terms of the two Higgs vacuum expectation values (VEV), v_1 and v_2 , and one physical Higgs mass (usually, $m_{H_3^0}$). Here, v_1 (v_2) is the vacuum expectation values of the Higgs field which couples exclusively to *down - type* (*up - type*) quarks and leptons. The value $v_1^2 + v_2^2$ is fixed by the W mass (or equivalently by the Fermi constant G_F),

$$v_1^2 + v_2^2 \approx (246 \text{ GeV})^2 \quad (3.1)$$

while the ratio v_2/v_1 is a free parameter of the model in terms of the angle β ;

$$\tan \beta = v_2/v_1 \quad (3.2)$$

The supersymmetric constraints imply that the MSSM Higgs sector is automatically $CP - conserving$ (at tree level). Thus, $\tan \beta$ is a real parameter (conventionally taken to be positive), and the physical neutral Higgs scalars are $CP - eigenstates$. Nevertheless, the MSSM does contain a number of possible new sources of $CP - violation$. For example, gaugino-mass parameters, the $A - parameters$, and μ may be complex. Some combination of these complex phases must be less than an order of $10^{-2} - 10^{-13}$ (for a

supersymmetric-breaking scale of 100 GeV) to avoid generating electric dipole moments for the neutron, electron and atoms in conflict with observed data. However, these complex phases have little impact on the direct searches for supersymmetric particles, and are usually ignored in experimental analysis.

3.2 Extended Higgs Sectors

The Higgs mechanism has solved the problem of having massive gauge bosons in an invariant local gauge theory without spoiling renormalizability and unitarity. This is achieved by means of a spontaneous breaking of the gauge symmetry in which the ground state (vacuum) loses part of the symmetry whereas the Lagrangian itself remains fully symmetric.

In the $SU(2)\times U(1)$ standard Glashow-Weinberg-Salam model (GWS or SM) the spontaneous symmetry breaking is induced by the presence of a doublet (under $SU(2)$) of complex scalar fields [12]

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}. \quad (3.3)$$

The new fields have Yukawa type interactions with matter fermion fields and also have self-interactions of the form

$$V(\phi) \equiv -\mu^2|\phi|^2 + \lambda|\phi|^4, \quad (3.4)$$

where μ^2 and λ are positive constants. After the Higgs mechanism, the theory contains - apart from the fields- 3 massive gauge bosons (W^+ , W^- , Z), 1 massless

photon and 1 physical scalar (H), the “Higgs boson”. The other three real scalars of the doublet (the “Goldstino bosons”) have become the longitudinal components of the three massive gauge bosons.

Although the minimal Higgs sector of the SM is sufficient to explain the generation of the fermion and gauge boson masses, more complicated structures in the scalar sector can not be excluded and are even unavoidable in many unifying extensions of the SM. These extended Higgs sectors have potentially richer phenomenology but are also subjected to phenomenological constraints, for example, the electroweak ρ –*parameter*, and the presence of tree level couplings of the type $W^-Z^0H^-$.

3.2.1 The ρ –Parameter constraint

The most important phenomenological constraint on the Higgs sector is the value of the electroweak ρ –parameter which, experimentally, is very close to 1

$$\rho \equiv \frac{m_W^2}{m_Z^2 \cos^2 \theta_W} \approx 1 \quad (3.5)$$

With an arbitrary Higgs sector consisting of several scalar multiplets ϕ_i of weak isospin $T_{(i)}$ and weak hypercharge $Y_{(i)}$, the ρ –parameter is given by

$$\rho = \frac{\sum_i \left[T_{(i)} (T_{(i)} + 1) - (Y_{(i)}/2)^2 \right] v_i^2 c_i}{2 \sum_i \left((Y_{(i)}/2)^2 \right) v_i^2 c_i} \quad (3.6)$$

where v_i is the VEV of the multiplet ϕ_i and $c_i = 1(\frac{1}{2})$ when $Y_{(i)} \neq 0$ ($= 0$).

It is easy to check that below $T = 10$ only the representations $(T, Y) = (0, 0), (\frac{1}{2}, \pm 1)$ and $(3, \pm 4)$ lead “naturally” (*i.e.* independently of the values of v_i) to $\rho = 1$. Even if we allow ρ to deviate from 1 by 1%, no new representations appear. Leaving aside the case of $(T, Y) = (3, \pm 4)$ involving scalars of electric charge $Q = 5$, only doublets and singlets are acceptable.

Of course other representations are allowed if we only require $\rho \approx 1$ for *some* values of the v_i . The simplest cases are the models with a doublets and a (real or complex) triplet but with ρ differing slightly from 1 unless $v_i = 0$. The simplest “unnatural” case with $\rho = 1$ is a model with one doublet and two triplets (one real and one complex) with equal VEV’s.

3.2.2 The $W^- Z^0 H^-$ Couplings

a general feature of the extended Higgs sectors is the presence of *physical* charged scalar fields ($H^\pm$). This fact implies a potentially rich phenomenology. In particular one expects a tree level couplings of the type $W^- Z^0 H^-$ in analogy to $W^- W^+ H$ and $Z^0 Z^0 H$ in the SM. However, it turns out that this coupling is absent (at the tree level) in the simplest “natural” extensions of the Higgs sector (*i.e.* with doublets and singlets) and it is small (proportional to $\sqrt{|1 - \rho|}$) in the simplest “unnatural” extensions. It is also easy to prove (using the electromagnetic gauge invariance) that the $W^- \gamma H^+$ couplings vanish at the tree level in all models.

3.3 The Two Higgs Doublet Model

A Higgs sector consisting of two scalar doublets

$$\phi_1 = \begin{pmatrix} \phi_1^+ \\ \phi_1^0 \end{pmatrix}, \quad \text{and} \quad \phi_2 = \begin{pmatrix} \phi_2^+ \\ \phi_2^0 \end{pmatrix} \quad (3.7)$$

is the simplest “natural” *extension* of the SM. As we have seen, in such case the $W^-Z^0H^+$ coupling is automatically absent at the tree level. The most general CP -conserving potential involving two doublets is [13]

$$\begin{aligned} V(\phi_1, \phi_2) = & \lambda_1(\phi_1^+\phi_1 - v_1^2)^2 + \lambda_2(\phi_2^+\phi_2 - v_2^2)^2 \\ & + \lambda_3 [(\phi_1^+\phi_1 - v_1^2)^2 + (\phi_2^+\phi_2 - v_2^2)^2]^2 \\ & + \lambda_4 [(\phi_1^+\phi_1)(\phi_2^+\phi_2) - (\phi_1^+\phi_2)(\phi_2^+\phi_1)] \\ & + \lambda_5 [\text{Re}(\phi_1^+\phi_2) - v_1v_2]^2 + \lambda_6 [\text{Im}(\phi_1^+\phi_2)]^2, \end{aligned} \quad (3.8)$$

where λ_i are 6 arbitrary real parameters. If $\lambda_i > 0$, the minimum of the potential corresponds to

$$\langle \phi_i \rangle = \begin{pmatrix} 0 \\ v_i \end{pmatrix}, \quad \text{and} \quad \langle \phi_2 \rangle = \begin{pmatrix} 0 \\ v_2 \end{pmatrix}. \quad (3.9)$$

In general the neutral components of the doublets can have flavor changing couplings to the fermions. These flavor changing neutral currents (FCNC) interactions can be suppressed (as required phenomenologically) either by giving large masses to these scalars or by arranging their Yukawa couplings to the fermions. *Glashow* and *Weinberg*

proved a theorem stating the sufficient condition for avoiding these FCNC effects; “(FCNC) interactions induced by neutral Higgs scalars are absent if all fermions of a given charge receive their masses from a single doublet”. This theorem is trivially satisfied in the SM since there is only one doublet available. It is also satisfied in the “Minimal supersymmetric model”.

Of the 8 available degrees of freedom (4 complex scalars with two real components), 3 are the Goldstone bosons to become the longitudinal components of the $W^\pm$ and Z^0 bosons that become massive by the “Higgs mechanism”. The remaining five scalars are *physical* states (two are charged and three neutral). They are

$$H^\pm = -\sin\beta\phi_1^\pm + \cos\beta\phi_2^\pm, \quad (3.10a)$$

$$H_1^0 = \sqrt{2}[(\text{Re}\phi_1^0 - v_1)\cos\alpha + (\text{Re}\phi_2^0 - v_2) - \sin\alpha], \quad (3.10b)$$

$$H_2^0 = \sqrt{2}[-(\text{Re}\phi_1^0 - v_1)\sin\alpha + (\text{Re}\phi_2^\pm - v_2) - \cos\alpha], \quad (3.10c)$$

$$H_3^0 = \sqrt{2}[\sin\beta\text{Im}\phi_1^0 + \cos\beta\text{Im}\phi_2^\pm], \quad (3.10d)$$

with masses $m_{H^\pm}$, $m_{H_1^0}$, $m_{H_2^0}$, and $m_{H_3^0}$, respectively. The angle α , and the masses are functions of the parameters λ_i and the VEV's $v_{1,2}$

$$m_{H^\pm}^2 = \lambda_4(v_1^2 + v_2^2), \quad (3.11a)$$

$$m_{H_3^0} = \lambda_6(v_1^2 + v_2^2), \quad (3.11b)$$

$$m_{H_1^0, H_2^0}^2 = \frac{1}{2} [A + C \pm D], \quad (3.11c)$$

$$\sin 2\alpha = \frac{2B}{D}, \quad (3.11d)$$

$$\cos 2\alpha = \frac{A - C}{D}, \quad (3.11e)$$

where

$$A = 4v_1^2(\lambda_2 + \lambda_3) + v_2^2\lambda_5, \quad (3.12a)$$

$$B = (4\lambda_3 + \lambda_5)v_1v_2, \quad (3.12b)$$

$$C = 4v_2^2(\lambda_2 + \lambda_3) + v_1^2\lambda_5, \quad (3.12c)$$

$$D = \sqrt{(A - C)^2 + 4B^2}. \quad (3.12d)$$

This model is completely specified by six parameters: $m_{H^\pm}$, $m_{H_1^0}$, $m_{H_2^0}$, $m_{H_3^0}$, α , and $\tan \beta (\equiv v_2/v_1)$.

In the absence of fermions, the Lagrangian (involving only gauge bosons and scalars) is C - and P -conserving and the gauge bosons have the following quantum numbers:

$$J^{PC} = 1^{--}(\gamma), \quad 1^{--}(Z), \quad \text{and} \quad J^P = 1^-(W). \quad (3.13)$$

Similarly, the physical scalars are fixed to be

$$J^{PC} = 0^{++}(H_1^0), 0^{++}(H_2^0), 0^{++}(H_3^0), \text{ and } J^P = 0^+(H^\pm). \quad (3.14)$$

As a consequence, the couplings $ZH_1^0H_1^0$ and $ZH_2^0H_2^0$ are zero since they would violate Bose symmetry. The coupling $ZH_1^0H_2^0$ also vanishes due to CP -conservation and couplings ZZH_3^0 and $W^-W^+H_3^0$ are forbidden by C -conservation. These last tree results hold to all orders of perturbation theory before fermions are introduced.

Other couplings are absent only at the tree level:

$\gamma H_i^0 H_j^0$, $\gamma\gamma H_i^0$, ggH_i^0 , $W^\pm\gamma H^\mp$, and $W^\pm ZH^\mp$, but they can be generated in higher orders of perturbation theory and can lead to interesting rare decays. All other couplings are in principle allowed. In particular, the couplings $W^-W^+H_{1,2}^0$ exist and satisfy the sum rule

$$g_{V^+V^-H_1^0}^2 + g_{V^+V^-H_2^0}^2 = g_{V^+V^-H^0}^2(SM), \quad V = (W, Z), \quad (3.15)$$

i.e. they are somewhat suppressed compared to the analogous SM couplings.

When fermions are introduced, since the couplings to fermions are not C - and P -conserving (although CP still approximately conserved), scalar and gauge bosons are regarded by fermions as mixtures of J^{PC} and $J^{(-P)(-C)}$ states. In particular, since a fermion-antifermion pair with zero total angular momentum always has $C = +$, in the $H_i^0 f\bar{f}$ couplings, the Higgs fields H_1^0 , H_2^0 , and H_3^0 acts respectively as 0^{++} , 0^{++} , and 0^{-+} states.

3.4 The Higgs Sector *of the* MSSM

One peculiar fact of all supersymmetric gauge theories is that at least two doublets are required. The simplest supersymmetric extension of the SM is the MSSM where the Higgs sector consists of just two doublets and it is a particular example of the two-doublet models. In this case, SUSY imposes constraints and the number of the independent parameters is reduced from six to two: $m_{H_3^0}$ and $\tan\beta$. The Higgs sector is then completely specified by the values of these two parameters.

The Higgs potential of the MSSM can be written as [14], [15]:

$$\begin{aligned}
V(\phi_1, \phi_2) = & m_1^2 \phi_1^\dagger \phi_1 + m_2^2 \phi_2^\dagger \phi_2 - m_{1,2}^2 (\phi_1^\dagger \phi_2 + \phi_2^\dagger \phi_1) \\
& + \frac{1}{8} (g'^2 + g^2) \left[(\phi_1^\dagger \phi_1)^2 + (\phi_2^\dagger \phi_2)^2 \right] \\
& + \frac{1}{4} (g'^2 - g^2) (\phi_1^\dagger \phi_1) (\phi_2^\dagger \phi_2) - \frac{1}{2} g^2 (\phi_1^\dagger \phi_2) (\phi_2^\dagger \phi_1),
\end{aligned} \tag{3.16}$$

where g' and g are the U(1) and SU(2) gauge couplings, respectively. Let V be broken spontaneously, then by comparing eq.(3.8) with eq.(3.16) we obtain the following results

$$\lambda_2 = \lambda_1 \quad (3.17a)$$

$$\lambda_3 = \frac{1}{8}(g^2 + g'^2) - \lambda_1 \quad (3.17b)$$

$$\lambda_4 = 2\lambda_1 - \frac{1}{2}g'^2 \quad (3.17c)$$

$$\lambda_5 = \lambda_6 = 2\lambda_1 - \frac{1}{2}(g^2 + g'^2) \quad (3.17d)$$

$$\lambda_7 = -\frac{1}{8}(v_1^2 - v_2^2)^2(g^2 + g'^2) \quad (3.17e)$$

$$m_1^2 = 2\lambda_1 v_2^2 - \frac{1}{4}(g^2 + g'^2)(v_1^2 + v_2^2) \quad (3.17f)$$

$$m_2^2 = 2\lambda_1 v_1^2 - \frac{1}{4}(g^2 + g'^2)(v_1^2 + v_2^2) \quad (3.17g)$$

$$m_{1,2}^2 = \frac{1}{2}v_1 v_2 (4\lambda_1 - g^2 - g'^2) \quad (3.17h)$$

Using eq. (3.17h) to eliminate λ_1 in eqs. (3.17f) and (3.17g) we get

$$m_1^2 = m_{1,2}^2 \frac{v_2}{v_1} - \frac{1}{4}(g^2 + g'^2)(v_1^2 + v_2^2) \quad (3.18)$$

$$m_2^2 = m_{1,2}^2 \frac{v_1}{v_2} - \frac{1}{4}(g^2 + g'^2)(v_1^2 + v_2^2) \quad (3.19)$$

Hence

$$m_1^2 + m_2^2 = m_{1,2}^2 (\tan \beta + \cot \beta) \quad (3.20)$$

$$v_1^2 + v_2^2 = \frac{-4m_1^2 \cos^2 \beta + 4m_2^2 \sin^2 \beta}{(g^2 + g'^2)(\cos^2 \beta - \sin^2 \beta)} \quad (3.21)$$

Using eqs. (3.12), and (3.17)-(3.21), we immediately get the spectrum of physical Higgs particles. The results are

$$\begin{aligned}
m_{H_1^0}^2 &= m_1^2 + m_2^2, \\
m_{H_2^0}^2 &= m_W^2 + m_{H_3^0}^2 \\
m_{H_1^0, H_2^0}^2 &= \frac{1}{2} \left[\sqrt{m_Z^2 + m_{H_3^0}^2 \pm (m_Z^2 + m_{H_3^0}^2)^2 - 4m_Z^2 m_{H_3^0}^2 \cos^2 2\beta} \right],
\end{aligned} \tag{3.22}$$

where

$$\begin{aligned}
m_W^2 &= \frac{1}{2} g^2 (v_1^2 + v_2^2), \\
m_Z^2 &= \frac{1}{2} (g^2 + g'^2) (v_1^2 + v_2^2)
\end{aligned} \tag{3.23}$$

are the squares of the masses of both W and Z bosons.

In this case the relations between α and β are

$$\begin{aligned}
\sin 2\alpha &= -\sin 2\beta \left(\frac{m_{H_1^0}^2 + m_{H_2^0}^2}{m_{H_1^0}^2 - m_{H_2^0}^2} \right), \\
\cos 2\alpha &= -\cos 2\beta \left(\frac{m_{H_3^0}^2 - m_Z^2}{m_{H_1^0}^2 - m_{H_2^0}^2} \right).
\end{aligned} \tag{3.24}$$

From these expressions, the following inequalities follow,

$$\begin{aligned}
m_W &< m_{H^\pm}, \\
m_{H_2^0} &< m_{H_3^0}, \\
m_{H_2^0} &< m_Z < m_{H_1^0}
\end{aligned}
\tag{3.25}$$

The fact that one of the neutral scalars, H_2^0 , is lighter than the Z boson, is an interesting results of the MSSM. The heavy scalar H_1^0 , on the other hand, has a $W^-W^+H_1^0$ coupling which is suppresses with respect to corresponding one in the SM by a factor

$$\left[\frac{m_{H_2^0}^2 (m_Z^2 - m_{H_2^0}^2)}{(m_{H_1^0}^2 + m_{H_2^0}^2) (m_{H_1^0}^2 + m_{H_2^0}^2 - m_Z^2)} \right]^2. \tag{3.26}$$

This factor decreases as $1/m_{H_1^0}^2$ where $m_{H_1^0}^2$ increases, and it is lower than ~ 0.15 for $m_{H_1^0}^2 > 2m_W$. The heavy scalar then behaves differently from the SM Higgs since the latter interacts more and more strongly with the W boson when m_H increases. This different behavior of SUSY theories is consistent with the fact that the ultraviolet cut-off of the latter is far above $\sim 1TeV$.

These two results, namely, the existence of a light Higgs boson and the decoupling of the heavy one from the gauge bosons are general results which survive in the more general supersymmetric models, including the “superstring inspired” ones.

Recent results from LEP experiments have restricted the allowed region in the $m_{H_1^0}$, $\tan \beta$ plane and on $m_{H^\pm}$.

To summarize, there are five physical Higgs particles in the MSSM, a charged Higgs pair ($H^\pm$), two CP -even neutral Higgs bosons (denoted by H_1^0 and H_2^0 where

$m_{H_1^0} > m_{H_2^0}$) and one CP -odd neutral Higgs boson¹ H_3^0 . The properties of the Higgs sector of the MSSM are determined by the Higgs potential which is made of quadratic terms and quartic interaction terms. The strength of the interaction terms are directly related to the gauge couplings by supersymmetry (and are not affected at tree-level by supersymmetry breaking). As a result, $\tan\beta$ and one Higgs mass ($m_{H_3^0}$) determine: The Higgs spectrum, an angle α (which indicates the amount of mixing of the original $Y = \pm 1$ Higgs doublet states in the physical CP -even scalars), and the Higgs boson couplings.

3.5 The Supersymmetric Particle Sector *of the* MSSM

The supersymmetric partner of the gauge and Higgs bosons are fermions, whose names are obtained by appending “*ino*” at the end of the SM particle name. The *gluino* is the color octet Majorana fermion partner of the gluon with mass $m_{\tilde{g}} = |M_3|$. The supersymmetric partner of the electroweak gauge and Higgs bosons (the *gauginos* and *Higgsinos*) can mix. As a result, the physical mass eigenstates are model-dependent linear combinations of these states, called *charginos* and *neutralinos*, which are obtained by diagonalizing the corresponding mass matrix [16].

The chargino-mass matrix depends on M_2 , μ , $\tan\beta$, and m_W . The chargino mass eigenstates are denoted by $\tilde{\chi}_1^+$, $\tilde{\chi}_2^+$ according to the convention that $\tilde{\chi}_1^+ \leq \tilde{\chi}_2^+$.

The neutralino mass matrix depends on M_1 , M_2 , μ , $\tan\beta$, m_Z , and the weak mixing

¹In recent reviews of particle properties, the symbol A^0 replaces H_3^0 , to denote the CP -odd neutral Higgs boson.

angle θ_W . The corresponding neutralino eigenstates are denoted by $\tilde{\chi}_i^0$ ($i = 1, \dots, 4$), according to the convention that $\tilde{\chi}_1^0 \leq \tilde{\chi}_2^0 \leq \tilde{\chi}_3^0 \leq \tilde{\chi}_4^0$.

If a chargino or a neutralino eigenstate approximates a particular gaugino or Higgsino state, it may be convenient to use the corresponding nomenclature. For example, if M_1 and M_2 are small compared to m_Z (and μ), then the lightest neutralino $\tilde{\chi}_1^0$ will be nearly a pure photino, $\tilde{\gamma}$ (the supersymmetric partner of the photon).

It is common to reduce the supersymmetric parameter freedom by requiring that all three gaugino-mass parameters are equal at some grand unification scale. Then, at the electroweak scale the gaugino-mass parameter can be expressed in terms of one of them which we choose to be $M_2 \equiv M$. The other two gaugino-mass parameters are given by

$$\begin{aligned} M_1 &= \frac{3}{5}M' = \left(\frac{g'^2}{g^2}\right) M, \\ M_3 &\equiv m_{\tilde{g}} = \left(\frac{g_s^2}{g^2}\right) M, \end{aligned} \tag{3.27}$$

where M' , M and $m_{\tilde{g}}$ are the bino masses respectively. Having made this assumption, the chargino and neutralino masses and mixing angles depend only on three unknown parameters: the *wino* mass M , the Higgs mass parameter μ , and $\tan\beta$.

The supersymmetric partners of the squarks and leptons are spin zero bosons: the *squarks*, charged *sleptons*, and *sneutrinos*.

3.5.1 The charginos

The charginos, $\tilde{\chi}_i^\pm$ ($i = 1, 2$), are four component Dirac fermions which arise due to mixing of winos, $\widetilde{W}^-$, $\widetilde{W}^+$ and the charged Higgsinos, $\widetilde{H}^-$, and $\widetilde{H}^+$ [17], [18]. Because there actually two independent mixings, $(\widetilde{W}^-, \widetilde{H}^-)$ and $(\widetilde{W}^+, \widetilde{H}^+)$, we shall need to define two unitary mixing matrices. We define in two components spinor notation:

$$(\psi_j^\pm)^T = \left(-i\widetilde{W}^\pm, \widetilde{H}^\pm \right), \quad \text{where } j = 1, 2 \quad (3.28)$$

The mass term in the Lagrangian is:

$$\mathfrak{L}_m = (\psi^-)^T \mathbf{X} \psi^+ + h.c., \quad (3.29)$$

where

$$\mathbf{X} = \begin{pmatrix} M & \sqrt{2}m_W \sin \beta \\ \sqrt{2}m_W \cos \beta & \mu \end{pmatrix}. \quad (3.30)$$

The mass matrix $\mathbf{X}$ is diagonalized by the unitary 2×2 matrices $\mathbf{U}$ and $\mathbf{V}$:

$$\mathbf{U}^* \mathbf{X} \mathbf{V}^{-1} = \mathbf{M}_D, \quad (3.31)$$

where $\mathbf{M}_D$ is the diagonal chargino mass matrix. In particular, $\mathbf{U}$ and $\mathbf{V}$ can be chosen so that the elements of the diagonal matrix $\mathbf{M}_D$ are real and *non-negative*. We define two component mass eigenstates via:

$$\begin{aligned}
\chi_1^+ &= V_{ij}\psi_j^+, \\
\chi_1^- &= V_{ij}\psi_j^-, \quad \text{where } i, j = 1, 2,
\end{aligned} \tag{3.32}$$

The proper four component mass eigenstates are the charginos which are defined in terms of two component mass eigenstates as:

$$\tilde{\chi}_1^+ = \begin{pmatrix} \chi_1^+ \\ \chi_1^- \end{pmatrix}, \quad \tilde{\chi}_2^+ = \begin{pmatrix} \chi_2^+ \\ \chi_2^- \end{pmatrix}, \tag{3.33}$$

The mass eigenvalues $\mathbf{M}_D$ (the two components of the diagonal) are given by

$$\mathbf{M}_D^2 = \frac{1}{2} \left\{ \begin{aligned} &|\mu^2| + |M^2| + 2m_W^2 \\ &\mp \sqrt{(|\mu^2| + |M^2| + 2m_W^2 - 4|\mu^2||M^2|)^2 - 4m_W^4 \sin^2 2\beta + 8m_W^2 \sin^2 2\beta \operatorname{Re}(\mu M)} \end{aligned} \right\} \tag{3.34}$$

If CP -violation effects are ignored (in such case, M and μ are real parameters), then one can choose a convention where $\tan \beta$ and M are positive- note that the relative sign of M and μ are meaningful. The sign of μ is convention dependent² Now eq. (3.34) becomes

$$\mathbf{M}_D^2 = \frac{1}{2} \left\{ \begin{aligned} &\mu^2 + M^2 + 2m_W^2 \\ &\pm \sqrt{(M^2 - \mu^2)^2 + 4m_W^4 \cos^2 2\beta + 4m_W^2 (M^2 + \mu^2 + 2M\mu \sin 2\beta)} \end{aligned} \right\} \tag{3.35}$$

²Notice that both sign conventions appear in literature.

and it has the roots

$$M_{D\,1,2} = \frac{1}{2} \left(\sqrt{(M - \mu)^2 + 2m_W^2 (1 + \sin 2\beta)} \mp \sqrt{(M + \mu)^2 + 2m_W^2 (1 - \sin 2\beta)} \right) \quad (3.36)$$

We write the chargino mass eigenvalues in the form $M_{D\,i} = \eta_i m_{\tilde{\chi}_i^\pm}$, $i, j = 1, 2$, with $m_{\tilde{\chi}_i^\pm} = |M_{D\,i}|$, and $\eta_i = \text{sign}(M_{D\,i}) = \pm 1$.

Assuming CP -conservation we choose the matrices $\mathbf{U}$ and $\mathbf{V}$ real. The matrix elements U_{ij} and V_{ij} are given by

$$U_{1,2} = U_{2,1} = \frac{\theta_1}{\sqrt{2}} \sqrt{1 + \frac{M^2 - \mu^2 - 2m_W^2 \cos 2\beta}{W}} \quad (3.37a)$$

$$U_{2,2} = -U_{1,1} = \frac{\theta_2}{\sqrt{2}} \sqrt{1 - \frac{M^2 - \mu^2 - 2m_W^2 \cos 2\beta}{W}} \quad (3.37b)$$

$$V_{2,1} = -V_{1,2} = \frac{\theta_3}{\sqrt{2}} \sqrt{1 + \frac{M^2 - \mu^2 + 2m_W^2 \cos 2\beta}{W}} \quad (3.37c)$$

$$V_{2,2} = V_{1,1} = \frac{\theta_4}{\sqrt{2}} \sqrt{1 - \frac{M^2 - \mu^2 + 2m_W^2 \cos 2\beta}{W}} \quad (3.37d)$$

Where the sign factors θ_i , $i = 1, \dots, 4$, are given in Table 3.1, and

$$W = \sqrt{(M^2 + \mu^2 + 2m_W^2)^2 - 4(M\mu - m_W^2 \sin 2\beta)^2} \quad (3.38)$$

θ_i	$\tan \beta > 1$	$\tan \beta < 1$
θ_1	1	ε_B
θ_2	ε_B	1
θ_3	ε_A	1
θ_4	1	ε_A

Table 3.1. Sign factors $\theta_i, i = 1, \dots$, where $\varepsilon_A = \text{sign}(M \sin \beta + \mu \cos \beta)$

and $\varepsilon_B = \text{sign}(M \cos \beta + \mu \sin \beta)$.

3.5.2 The Neutralinos

The neutralinos, $\tilde{\chi}_i^0 (i = 1, \dots, 4)$, are four-component Majorana fermions which arise due to mixing of the two neutral gauginos $\tilde{B}$ (Bino), $\tilde{W}^3$ (neutral W -ino), and the two neutral Higgsinos, $\tilde{H}_1^0$, and $\tilde{H}_2^0$ [19],[20],[21]. As basis of the neutral gaugino-Higgsino system we conveniently take

$$(\psi^0)^T = (-i\tilde{B}, -i\tilde{W}^3, \tilde{H}_1^0, \tilde{H}_2^0), \quad (3.39)$$

The mass term in the Lagrangian is:

$$\mathfrak{L}_m = \frac{1}{2} (\psi^0)^T \mathbf{Y} \psi^0 + h.c., \quad (3.40)$$

where

$$\mathbf{Y} = \begin{pmatrix} M' & 0 & -m_Z \sin \theta_W \cos \beta & -m_Z \sin \theta_W \cos \beta \\ 0 & M & m_Z \cos \theta_W \sin \beta & -m_Z \cos \theta_W \sin \beta \\ -m_Z \sin \theta_W \cos \beta & m_Z \cos \theta_W \cos \beta & 0 & -\mu \\ m_Z \sin \theta_W \sin \beta & -m_Z \cos \theta_W \sin \beta & -\mu^2 & 0 \end{pmatrix} \quad (3.41)$$

on the $(-i\tilde{B}, -i\tilde{W}^3, \tilde{H}_1^0, \tilde{H}_2^0)$ basis. The two-component mass eigenstates can be obtained by diagonalizing the mass matrix $\mathbf{Y}$

$$\tilde{\chi}_i^0 = N_{ij} \psi_j^0, \quad (3.42)$$

where N is a complex unitary matrix ($\mathbf{N}^\dagger \mathbf{N} = \mathbf{1}$) satisfying

$$\mathbf{N}^* \mathbf{Y} \mathbf{N}^{-1} = \mathbf{N}_D, \quad (3.43)$$

and $\mathbf{N}_D$ is the diagonal neutralino mass matrix. the four-component mass eigenstates are the neutralinos which are defined in terms of two-component mass eigenstates:

$$\tilde{\chi}_i^0 = \begin{pmatrix} \chi_i^0 \\ \bar{\chi}_i^0 \end{pmatrix}, \quad i = 1, \dots, 4. \quad (3.44)$$

Assuming CP -invariance $\mathbf{N}$ is replaced by another matrix $\mathbf{Z}$. This implies the changing of eq.(3.43) to

$$\mathbf{Z}^* \mathbf{Y} \mathbf{Z}^{-1} = \mathbf{M}_D, \quad (3.45)$$

and we write the neutralino mass eigenvalues in the form $M_{D\ i} = \varepsilon_i m_{\tilde{\chi}_i^0}$, $i = 1, \dots, 4$, with $m_{\tilde{\chi}_i^0} = |M_{D\ i}|$, and $\varepsilon_i = \text{sign}(M_{D\ i}) = \pm 1$. The relation between the $\mathbf{N}$ and $\mathbf{Z}$ matrices is

$$N_{ij} = \sqrt{\varepsilon_i} Z_{ij}, \quad (\text{no sum over } i). \quad (3.46)$$

Using the theory of equations, the expressions for the $M_{D\ i}$ (the four components of the diagonal) are given by

$$M_{D\ 1} = -A + B_+ + \frac{1}{4}C_+, \quad (3.47a)$$

$$M_{D\ 2} = +A - B_- + \frac{1}{4}C_+, \quad (3.47b)$$

$$M_{D\ 3} = -A - B_+ + \frac{1}{4}C_+, \quad (3.47c)$$

$$M_{D\ 4} = +A + B_- + \frac{1}{4}C_+, \quad (3.47d)$$

where

$$A = \sqrt{\frac{1}{2}a - \frac{1}{6}b}, \quad (3.48a)$$

$$B_{\pm} = \sqrt{-\frac{1}{2}a - \frac{1}{3}b \pm \frac{c}{\sqrt{8a - \frac{8}{3}b}}}, \quad (3.48b)$$

$$C_{\pm} = M' \pm M, \quad (3.48c)$$

and

$$a = \frac{1}{\sqrt[3]{2}} \operatorname{Re} \left(c^2 + \frac{2}{27}b^3 - \frac{8}{3}bd + i\sqrt{\frac{D}{27}} \right)^{\frac{1}{3}}, \quad (3.49a)$$

$$b = E - \frac{3}{8}C_+^2, \quad (3.49b)$$

$$c = \frac{1}{8}C_+^3 + \frac{1}{2}C_+E + C_+\mu^2 + Fm_Z^2 - \mu m_Z^2 \sin 2\beta, \quad (3.49c)$$

$$d = F\mu m_Z^2 \sin 2\beta - M'M\mu^2 + \frac{1}{16}EC_+^2 - \frac{3}{256}C_+^4 + \frac{1}{4}C_+(C_+\mu^2 + Fm_Z^2 - \mu m_Z^2 \sin 2\beta), \quad (3.49d)$$

where

$$D = -4 \left(-\frac{1}{3}b^3 - 4d \right)^3 - 27 \left(-c^2 - \frac{2}{27}b^3 + \frac{8}{3}bd \right)^2, \quad (3.50a)$$

$$E = M'M - m_Z^2 - \mu^2, \quad (3.50b)$$

$$F = M' \cos^2 \theta_W + M \sin^2 \theta_W. \quad (3.50c)$$

The elements of the mixing matrix $\mathbf{Z}$ are given by

$$Z_{i1} = \frac{1}{\sqrt{1 + G_i^2 + H_i^2 + I_i^2}} \quad (3.51a)$$

$$Z_{i2} = G_i Z_{i1}, \quad (3.51b)$$

$$Z_{i3} = H_i Z_{i1}, \quad (3.51c)$$

$$Z_{i4} = I_i Z_{i1}, \quad i = 1, \dots, 4, \quad (3.51d)$$

where

$$G_i = -\frac{J'_i}{J_i \tan \theta_W}, \quad (3.52a)$$

$$H_i = \frac{\mu J_i J'_i - \frac{1}{2} K_i}{L_i}, \quad (3.52b)$$

$$I_i = \frac{-M_{D\ i} J_i J'_i - K_i}{L_i}, \quad (3.52c)$$

$$J_i = M - M_{D\ i}, \quad J'_1 = M' - M_{D\ i}, \quad (3.52d)$$

$$K_i = m_Z^2 \sin 2\beta (C_- \cos^2 \theta_W + J_i), \quad (3.52e)$$

$$L_i = m_Z J_i \sin \theta_i (\mu \cos \beta + M_{D\ i} \sin \beta). \quad (3.52f)$$

3.5.3 The Sferminos

For a given fermion f , there are two supersymmetric partners $\tilde{f}_L$ and $\tilde{f}_R$ (sfermions) which are scalar parameters of the corresponding left and right-handed fermion. There are no $\tilde{\nu}_R$. However, in general, $\tilde{f}_L$ and $\tilde{f}_R$ are not mass-eigenstates since there is $\tilde{f}_L - \tilde{f}_R$ mixing which is proportional in strength to the corresponding element of the scalar mass-squared matrix [22]:

$$M_{LR}^2 = \begin{cases} m_d (A_d - \mu \tan \beta), & \text{for "down"-type } f \\ m_u (A_u - \mu \cot \beta), & \text{for "up" type } f \end{cases}, \quad (3.53)$$

where $m_d (m_u)$ is the mass of the appropriate "down" ("up") type quark or lepton.

Here, A_d and A_u are (unknown) soft-supersymmetric-breaking A -parameters and μ and $\tan \beta$ have been defined earlier. The signs of the A -parameters are also convenient-dependent due to the appearance of the *fermion* mass in eq.(3.53), one expects M_{LR} to be small compared to the diagonal squark and slepton masses, with the possible

exception of the top-squark, since m_t is large, and the bottom-squark and tau-slepton if $\tan \beta \gg 1$.

The (diagonal) L – and R – type squark and slepton masses are given by

$$m_{\tilde{u}_L}^2 = M_{\tilde{Q}}^2 + m_u^2 + m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \right), \quad (3.54a)$$

$$m_{\tilde{u}_R}^2 = M_{\tilde{U}}^2 + m_u^2 + \frac{2}{3} m_Z^2 \cos 2\beta \sin^2 \theta_W, \quad (3.54b)$$

$$m_{\tilde{d}_L}^2 = M_{\tilde{Q}}^2 + m_d^2 - m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{1}{3} \sin^2 \theta_W \right), \quad (3.54c)$$

$$m_{\tilde{d}_R}^2 = M_{\tilde{D}}^2 + m_d^2 - \frac{1}{3} m_Z^2 \cos 2\beta \sin^2 \theta_W, \quad (3.54d)$$

$$m_{\tilde{\nu}}^2 = M_{\tilde{L}}^2 + \frac{1}{3} m_Z^2 \cos 2\beta, \quad (3.54e)$$

$$m_{\tilde{e}_L}^2 = M_{\tilde{L}}^2 + m_e^2 - m_Z^2 \cos 2\beta \left(\frac{1}{2} - \sin^2 \theta_W \right), \quad (3.54f)$$

$$m_{\tilde{e}_R}^2 = M_{\tilde{E}}^2 + m_e^2 - m_Z^2 \cos 2\beta \sin^2 \theta_W, \quad (3.54g)$$

The soft-supersymmetry-breaking parameters: $M_{\tilde{Q}}^2, M_{\tilde{U}}^2, M_{\tilde{D}}^2, M_{\tilde{L}}^2, M_{\tilde{E}}^2$ are unknown parameters. In the equations above, the notation of the first generation fermions has been used and generational indices have been suppressed. further complications such as integrational mixing are possible, although there are some constraints from the non-observation of flavor-changing current (FCNC).

3.6 Reducing *the* MSSM Parameter Freedom

One way to guarantee the absence of FCNC's mediated by virtual supersymmetric-particle exchange is to posit that the diagonal soft-supersymmetry-breaking scalar squared-

masses are universal in flavor space at some energy scale (normally taken to be at or near the Plank scale). Renormalization group evolution is used to determine the low energy values for the scalar mass parameters listed above. This assumption reduces the MSSM parameter freedom. For example, the supersymmetric grand unified models with universal scalar masses at Plank scale typically give $M_{\tilde{L}} \approx M_{\tilde{E}} < M_{\tilde{Q}} \approx M_{\tilde{U}} \approx M_{\tilde{D}}$ with the squark masses somewhere between a factor of 1-3 larger than the slepton masses (neglecting generational distinction). More specifically, the first two generations are thought to be degenerate in mass, while $M_{\tilde{Q}_3}$ and $M_{\tilde{U}_3}$ are typically reduced by a factor of 1-3 from the other soft supersymmetric breaking masses because of renormalization effects due to the heavy top quark masses.

As a result, four flavors of the squarks (with two squarks eigenstates per flavor) and $\tilde{b}_R$ will be nearly mass-degenerate and somewhat heavier than six flavors of nearly mass-degenerate sleptons (with two per flavor for the charged sleptons and one for the sneutrinos). On the other hand, $\tilde{b}_L$ mass and the diagonal $\tilde{t}_L$ and $\tilde{t}_R$ masses are reduced compared to the common squark mass of the first two generations. In addition, third generation squark masses and tau-slepton masses are sensitive to the respective $\tilde{f}_L - \tilde{f}_R$ mixing as discussed before.

Two additional theoretical frameworks are often introduced to reduce further the MSSM parameter freedom. the first involves grand unified theories (GUTs) and the desert hypothesis (*i.e.* no new physics between the TeV-scale and the GUT-scale). Perhaps one of the most compelling hints for low energy supersymmetry is the unification

of $SU(3) \times SU(2) \times U(1)$ gauge coupling predicted by supersymmetric GUT models (with the supersymmetry breaking scale of order 1 TeV or below).

The unification, which takes place at an energy scale of order $10^{16} GeV$ is quite robust (and depends weakly on the details of the GUT-scale theory). For example, a recent analysis finds that supersymmetric GUT unification implies that $\alpha_s(m_Z) = 0.129 \pm 0.010$, not including threshold corrections due to GUT-scale particles (which could diminish the value of $\alpha_s(m_Z)$). This result is compatible with the world average of $\alpha_s(m_Z) = 0.118 \pm 0.003$. In contrast, gauge coupling unification in the simplest non-supersymmetric GUT models fails by many standard deviations.

3.6.1 Minimization of the Higgs potential

In the MSSM, the Higgs sector has two unknown parameters, usually taken to be $\tan \beta \equiv v_2/v_1$ and $m_{H_3^0}$, the mass of its one physical pseudoscalar particle. Numerous phenomenological studies have been made using these parameters as continuous variables. However, there is an argument for $m_{H_3^0} = m_Z$ at the tree level and perhaps also $\tan \beta > \sqrt{3}$, by minimizing the minimum of the Higgs potential along a certain direction in parameter space [23].

The part of V (see eq.(316)) involving only neutral fields depends on four parameters: m_1^2 , m_2^2 , $m_{1,2}^2$, and $g_1^2 + g_2^2$. At its minimum V_0 , we can choose to keep m_1^2 and m_2^2 , but replace $m_{1,2}^2$ with $\tan \beta$ through eq. (3.20) and $g_1^2 + g_2^2$ with $v_1^2 + v_2^2$ through eq.(3.21). In that case,

$$V_0 = \frac{1}{2}(v_1^2 + v_2^2) (\cos^2 \beta - \sin^2 \beta)(m_1^2 \cos^2 \beta - m_2^2 \sin^2 \beta) \quad (3.55)$$

We now seek to minimize V_0 in parameter space. This is based on the assumption that the dynamic mechanism responsible for the soft breaking of supersymmetry may be such that the lowest possible value of V_0 is automatically chosen. It is also clear that $v_1^2 + v_2^2$, m_1^2 , m_2^2 set the energy scale of the symmetry breaking and V_0 has no lower bound as a function of these parameters. We should therefore consider them as fixed and vary $\sin^2 \beta$ to minimize V_0 . Let $x \equiv \sin^2 \beta$, then

$$\frac{\partial V_0}{\partial x} = \frac{1}{2} (v_1^2 + v_2^2) [- (3m_1^2 + m_2^2) + 4 (m_1^2 + m_2^2) x], \quad (3.56)$$

and

$$\frac{\partial^2 V_0}{\partial x^2} = 2 (v_1^2 + v_2^2) (m_1^2 + m_2^2). \quad (3.57)$$

Hence, the minimization of V_0 is achieved if

$$x = \frac{3m_1^2 + m_2^2}{4(m_1^2 + m_2^2)} \quad (3.58)$$

and $m_1^2 + m_2^2 > 0$, which is consistent with eq.(3.22). Using eq. (3.21) we then find

$$m_1^2 + m_2^2 = \frac{1}{2} (g_1^2 + g_2^2) (v_1^2 + v_2^2), \quad (3.59)$$

or equivalently,

$$m_{H_3^0} = m_Z. \quad (3.60)$$

This implies

$$m_{H^\pm} = \sqrt{m_W^2 + m_Z^2} \approx 121 \text{ GeV} \quad (3.61)$$

and

$$m_{H_1^0, H_2^0} = m_Z \sqrt{1 \pm \sin 2\beta} \quad (3.62)$$

From eq.(3.58), we find

$$\tan^2 \beta = \frac{3m_1^2 + m_2^2}{m_1^2 + 3m_2^2}, \quad (3.63)$$

which shows that if $m_1^2 > 0$ and $m_2^2 > 0$, then $\frac{1}{\sqrt{3}} < \tan \beta < \sqrt{3}$. However, because ϕ_2 couples to the *top* quark with a large Yukawa coupling, m_2^2 is expected to differ from m_1^2 by a large negative contribution from the renormalization-group equations, hence the case $m_1^2 > 0$ and $m_2^2 < 0$ should be considered. We then obtain

$$\tan \beta > \sqrt{3} \approx 1.731, \quad (3.64)$$

where $m_1^2 > 3|m_2^2|$ has also been assumed or else V_0 would have been minimized at $\sin^2 \beta > 1$, which is impossible. Using eq.(3.62) we find $m_{H_1^0} > 33 \text{ GeV}$ and $m_{H_2^0} < 125 \text{ GeV}$, with the constraints that $m_{H_1^0}^2 + m_{H_2^0}^2 = 2m_Z^2$. Experimentally, there is no evidence for the existence of any of the five scalar particles of the MSSM from the Z decay or in any other process.

3.6.2 Diophantine Analysis *of the* Higgs Boson

Diophantine quantization involves treating mass relations as Diophantine equations and seeking solutions in integers, analogous to the sets $(3, 4, 5), (5, 12, 13), etc.$, for the Pythagorean equation $x^2 + y^2 = z^2$. It was first applied to the Gell-Mann-Okubo meson-mass relation [24]

$$m_\pi^2 + 3m_\eta^2 = 4m_K^2, \quad (3.65)$$

for which the simplest nontrivial solution was the set $(2, 8, 7)$. The procedure not only gave integers proportional to the experimental masses $m_\pi = 135 - 140 MeV$, $m_\eta = 547 MeV$, $m_K = 494 - 498 MeV$ but also set a unit mass- the GCF of the three mesons masses- of the $70 MeV$ $(= (\hbar c/e^2) m_e c^2)$, as originally proposed by Nambu.

There is an analogy to this in the standard model. If one looks carefully at the values of the W and Z masses

$$m_W = 80.22 \pm 0.26 GeV, \quad (3.66)$$

$$m_Z = 91.173 \pm 0.020 GeV, \quad (3.67)$$

one finds that within experimental limits,

$$\cos \theta_W = m_W/m_Z = 15/17. \quad (3.68)$$

not only is this the ratio of the two integers, but these two particular two integers (15, 17) happen to be two sides of a right-angled triangle, then θ_W is, quite interestingly, a rational angle. A simple consequence of this, which we shall use below, is that the sine and cosine of such an angle must be rational.

A lengthy Diophantine analysis [25] applied to the scalar Higgs gauge boson mass relations of eq.(3.22) of minimal supersymmetry results in the following relations

$$\sin \beta = \begin{cases} \frac{8}{17}, & \beta < \pi/4 \\ \frac{15}{17}, & \beta > \pi/4 \end{cases}, \quad (3.69)$$

$$m_{H_3^0} = m_Z = 91.2 GeV, \quad (3.70a)$$

$$m_{H_1^0} = 37.5 GeV, \quad (3.70b)$$

$$m_{H_2^0} = 123 GeV, \quad (3.70c)$$

$$m_{H^\pm} = 121.5 GeV. \quad (3.70d)$$

Furthermore, most surprising of all, if we recall that the Weinberg angle θ_W is characterized by eq.(3.68), we have a relation between Higgs and Weinberg angles

$$\begin{cases} \theta_W, & \beta < \pi/4 \\ \frac{\pi}{4} - \theta_W, & \beta > \pi/4. \end{cases} \quad (3.71)$$

Finally, if $\beta > \pi/4$, $\sin \beta = \frac{15}{17}$, where the infrared quasi-fixed-point solution yields a *top*-quark mass

$$m_t \cong (190 - 210) \sin \beta \text{GeV} \approx 168 - 175 \text{GeV} \quad (3.72)$$

within the range of recent data [26].

3.7 Conclusion

The minimal supersymmetric standard model (*MSSM*) is the simplest extension of the standard model (*SM*) that includes soft broken supersymmetry (*SUSY*). The word “minimal” refers to minimal content of particles and gauge groups.

The *MSSM* contains a great number of free parameters which considerably limit the predictive power of the model. These are some commonly used ways to reduce the number of free constants and parameters in this theory. The most often employed method is to obtain values of parameters at the scale of the order of m_W by renormalization-group equations from the coupling constants of the supergravity theories investigated at the Plank mass. usually such theories are much more unified and contain typically only few free numbers. Of course, there are also many constraints originating from the experimental data-first of all the masses of the superpartners are bounded from below by their absence in the present experiments, but one can find also many more subtle limits.

The usual *MSSM* framework assumptions are:

1. *R*-parity is conserved. this assumption is commonly made in supersymmetry studies, as it prevents protons from decaying too quickly.
2. The lightest supersymmetric particle (LSP) is the lightest neutralino.

3. The intergenerational mixing in the squark, slepton, and quark sectors is small and can be neglected.
4. The four *left-handed* squarks of the first two generations are nearly degenerate at low energy with mass $m_{\tilde{q}}$ as are all six left-handed sleptons with mass $m_{\tilde{l}}$:

$$\begin{aligned}
m_{\tilde{u}_L} &\approx m_{\tilde{d}_L} \approx m_{\tilde{c}_L} \approx m_{\tilde{s}_L} \approx m_{\tilde{q}_L}, \\
m_{\tilde{\nu}_{e_L}} &\approx m_{\tilde{e}_L} \approx m_{\tilde{\nu}_{\mu_L}} \approx m_{\tilde{\mu}_L} \approx m_{\tilde{\nu}_{\tau_L}} \approx m_{\tilde{\tau}_L} \approx m_{\tilde{l}}.
\end{aligned} \tag{3.73}$$

5. The gaugino mass M_i , the parameter μ and $\tan\beta$ may be taken to be real, so that *CP – violation* plays no role.

Unification of gaugino mass, *i.e.* M_1 and M_2 are not independent.

Part II

Phenomenological Calculations

CHAPTER 4

INTRODUCTION

The phenomenological predictions of supersymmetry (SUSY) may be divided into three categories [27], [28], [29]:

1. Reflections of supersymmetric lagrangian in Standard Model (SM) phenomenology, including relations among the gauge coupling constants from supersymmetric grand unification and the presence of a heavy top quark and light Higgs scalar;
2. The prediction of new particles with the correct spin and quantum numbers assignments to be superpartners of the standard model particles; and
3. Well-defined quantitative relations among the couplings and masses of these new particles.

While the predictions of (1) are of great interest, their verification is clearly no substitute for direct evidence. The discovery of a large number of particles in category (2) would be strong support for supersymmetry. On the other hand, the most compelling confirmation of supersymmetry would likely be the precise verification of the relations of category (3). This would be specially true if, initially, only a small set of candidate supersymmetric partners are observed.

Most discussions of supersymmetry at future high energy colliders have concentrated on the question of particle searches. From one point of view, this is reasonable, because the existence of supersymmetric partners is unproven and this is a prerequisite for any further analysis. On the other hand, the discovery of the first evidence for supersymmetry, or for any other theoretical extension of the standard model, will begin a program of detailed experimental investigation of the new sector of particles required by this extension.

Supersymmetry provides a particularly interesting subject for studies of the detailed analysis of physics beyond the standard model. Supersymmetry are weakly coupled, and so their consequences can be worked out straightforwardly using perturbative computations. At the same time, supersymmetric models depend on a large number of unknown parameters, and different choices for these parameters yield qualitatively different realizations of possible new physics. Thus the phenomenology of supersymmetry is quite complex. Eventually, if supersymmetry does give a correct model of nature, the colliders of the next generation will be expected to determine the supersymmetric parameters, and their values will become clues that take us a step closer to a fundamental theory.

In the minimal supersymmetric extension of the standard model, MSSM, among the lightest supersymmetric particles there are four neutralinos (the supersymmetric partners of the neural electroweak gauge and Higgs bosons). In most scenarios, apart from the lightest supersymmetric particle (LSP), which is in general assumed to be the lightest neutralino ($\tilde{\chi}_1^0$) (stable and invisible), the particles that could be first observed

at future colliders are the next-to-lightest neutralino ($\tilde{\chi}_2^o$) and the light chargino ($\tilde{\chi}_1^\pm$) [30]. Therefore, any reasonable large supersymmetric signal must involve either the second lightest neutralino $\tilde{\chi}_2^o$ or the lighter charginos $\tilde{\chi}_1^\pm$. In general, we can not assume that the second lightest neutralino is heavier than the lighter chargino, since, $m_{\tilde{\chi}_2^o}$ is not independent of $m_{\tilde{\chi}_1^o}$ and $m_{\tilde{\chi}_1^\pm}$. In fact, in the region of parameter space in which charginos production is accessible to the future e^-e^+ colliders, $m_{\tilde{\chi}_2^o}$ and $m_{\tilde{\chi}_1^\pm}$ are very roughly degenerate, with the mass difference typically in the range

$$-10\text{GeV} \leq m_{\tilde{\chi}_2^o} - m_{\tilde{\chi}_1^\pm} \leq 20\text{GeV}.$$

When $m_{\tilde{\chi}_2^o} < m_{\tilde{\chi}_1^\pm}$, it is possible for the lighter chargino to decay through a cascade decays to a $\tilde{\chi}_2^o$, which in turn decays to an LSP.

The e^-e^+ colliders have been playing complementary roles to the hadron colliders in supersymmetry searches. In general, e^-e^+ colliders have reasonable signal rates in a clean environment with a definite center-of-mass energy, enabling us to perform precision measurements of particles' masses, lifetimes, and various different cross-sections, while hadron colliders provide opportunities to quickly survey high energy frontier. In particular, the production of $\tilde{\chi}_1^o \tilde{\chi}_2^o$ pairs at e^-e^+ colliders could allow the study of a wide region of the supersymmetry parameter space.

Owing to the relatively large cross-section of two body final state reactions, they can be used to search for supersymmetric particles with masses up to the beam energy. In this study, the production of certain three body final state reactions are calculated

to improve the sensitivity in searching for supersymmetric particles.

Outline of Part Two

In this part "the calculation part", reactions' cross sections are calculated and the corresponding curves are plotted with the results tabulated for a convenient reference.

Work went as follows:

1. In chapter five, the reaction $e^-e^+ \rightarrow H^-\tilde{\chi}_1^+\tilde{\chi}_1^o$ is considered.
2. In chapter six, the reaction $e^-e^+ \rightarrow h\tilde{\chi}_1^+\tilde{\chi}_1^-$ is considered.
3. In chapter seven, the reaction $e^-e^+ \rightarrow h\tilde{\chi}_1^o\tilde{\chi}_1^o$ is considered.
4. In chapter eight, the reaction $e^-e^+ \rightarrow hH^+H^-$ is considered.

CHAPTER 5

PRODUCTION OF A CHARGED HIGGS BOSON WITH A CHARGINO AND A NEUTRALINO

5.1 Introduction

In this chapter, the total cross section for the process $e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^o(p5)$ for different topologies and propagators (see Appendix A) are calculated and represented graphically. There are 28 different Feynman diagrams (tree level approximation) for which we gave the matrix element corresponding to each diagram. Diagrams with the same topology that can be obtained by changing the indices were represented once.

Work will go as follows:

1. Feynman diagrams are given,
2. Diagrams with the same topology are represented once, but has been taken into considerations when calculating the cross section.
3. Matrix elements are written, all the four momenta squares are defined to be mass squared (> 0),
4. Matrix elements are squared,

5. An average over the initial spin polarizations of the electron and positron pair and a sum over the final spin states of the outgoing particles arising from each initial spin state is carried out.
6. Results are represented graphically, and summarized in subsequent tables.

5.2 Feynman Diagrams

The following is the set of Feynman diagrams which were used to calculate the cross section of the associated production of a charged Higgs boson with a chargino and a neutralino. Our momentum notation is: $e^-(p_1)$, $e^+(p_2)$, $H^-(p_3)$, $\tilde{\chi}_1^+(p_4)$ and $\tilde{\chi}_1^0(p_5)$.




Figure 5.1: Feynman diagrams for the reaction: $e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^0(p5)$




Figure 5.2: Cont. Feynman diagrams for the reaction: $e^-(p_1)e^+(p_2) \rightarrow H^-(p_3)\tilde{\chi}_1^+(p_4)\tilde{\chi}_1^0(p_5)$

5.3 Matrix Elements

The following is the set of matrix elements corresponding to feynman diagrams in figures 5.1 and 5.2 used in our calculations:

$$\begin{aligned}\mathcal{M}_1 &= \bar{v}(p_2)A\gamma_\mu u(p_1)P_\gamma^{\mu\nu}(p_1+p_2)J(p_3+p_4-p_5)_\nu D_{H^+}(p_3+p_4)\bar{u}(p_4) \\ &\quad (Q_{ij}^L P_L + Q_{ij}^R P_R)v(p_3)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_2 &= \bar{v}(p_2)A\gamma_\mu u(p_1)P_\gamma^{\mu\nu}(p_1+p_2)\bar{u}(p_4)V\gamma_\nu D_{\bar{\chi}_1^+}(p_3+p_4)(\not{p}_3 + \not{p}_4 + m_{\bar{\chi}_1^+}) \\ &\quad (Q_{ij}^L P_L + Q_{ij}^R P_R)v(p_3)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_3 &= \bar{v}(p_2)\gamma_\mu(B^L P_L + B^R P_R)u(p_1)P_Z^{\mu\nu}(p_1+p_2)H(p_1+p_2)_\nu D_{H^+}(p_3+p_4) \\ &\quad \bar{u}(p_4)(Q_{ij}^L P_L + Q_{ij}^R P_R)v(p_3)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_{4,5} &= \bar{v}(p_2)\gamma_\mu(B^L P_L + B^R P_R)u(p_1)P_Z^{\mu\nu}(p_1+p_2)\bar{u}(p_4)\gamma_\nu(W_{ij}^L P_L + W_{ij}^R P_R) \\ &\quad D_{\bar{\chi}_k^+}(p_3+p_4)(\not{p}_3 + \not{p}_4 + m_{\bar{\chi}_k^+})(Q_{ij}^L P_L + Q_{ij}^R P_R)v(p_3)\end{aligned}$$

Where $k = 1, 2$.

$$\begin{aligned}\mathcal{M}_{6,7,8,9} &= \bar{v}(p_2)\gamma_\mu(B^L P_L + B^R P_R)u(p_1)P_Z^{\mu\nu}(p_1+p_2)\bar{u}(p_3)\gamma_\nu(S_{ij} + S'_{ij}\gamma_5) \\ &\quad D_{\bar{\chi}_k^0}(p_4+p_5)(\not{p}_4 + \not{p}_5 + m_{\bar{\chi}_k^0})(Q_{ij}^L P_L + Q_{ij}^R P_R)\bar{u}(p_4)\end{aligned}$$

Where $k = 1, 2, 3, 4$.

$$\begin{aligned} \mathcal{M}_{10,16,17,18,19,24,25,26} &= \bar{v}(p_2)(N_{hi} + N'_{hi}\gamma_5)v(p_3)D_{\tilde{e}_h}(p_1 - p_3)\bar{u}(p_3)(Q_{ij}^L P_L + Q_{ij}^R P_R) \\ &\quad D_{\tilde{\chi}_k^o}(p_3 + p_5)(\not{p}'_3 + \not{p}'_5 + m_{\tilde{\chi}_k^o})(N_{hi} - N'_{hi}\gamma_5)u(p_1) \end{aligned}$$

Where $k = 1, 2, 3, 4$.

and $h = L, R$.

$$\begin{aligned} \mathcal{M}_{11,12,13,14,20,21,22,23} &= \bar{v}(p_2)(N_{hi} + N'_{hi}\gamma_5)(\not{p}'_4 + \not{p}'_5 + m_{\tilde{\chi}_k^o})D_{\tilde{\chi}_k^o}(p_4 + p_5)\bar{u}(p_4)(Q_{ij}^L P_L + Q_{ij}^R P_R) \\ &\quad D_{\tilde{e}_h}(p_1 - p_3)\bar{u}(p_3)(N_{hi} - N'_{hi}\gamma_5)u(p_1) \end{aligned}$$

again, $k = 1, 2, 3, 4$.

and $h = L, R$.

$$\begin{aligned} \mathcal{M}_{27,28} &= \bar{v}(p_2)(T_i^L P_L + T_i^R P_R)v(p_4)D_{\tilde{\nu}_e}(p_2 - p_4)\bar{u}(p_3)(Q_{ij}^L P_L + Q_{ij}^R P_R)(\not{p}'_3 + \not{p}'_5 + m_{\tilde{\chi}_k^+}) \\ &\quad D_{\tilde{\chi}_k^+}(p_3 + p_5)(T_i^L P_L + T_i^R P_R)u(p_1) \end{aligned}$$

here $k = 1, 2$.

$$\begin{aligned} \mathcal{M}_8 &= \bar{v}(p_2)T_i^R P_R v(p_4)D_{\tilde{e}_L}(p_2 - p_4)L(p_1 + p_2 - p_3 - p_4)D_{\tilde{\nu}_e}(p_1 - p_3) \\ &\quad \bar{v}(p_3)(T_i^L P_L + T_i^R P_R)u(p_1) \end{aligned}$$

Where:

$$D_X(q) = \frac{1}{(q^2 - m_X^2)},$$

$$P_{Z,\gamma}^{\mu\nu} = \frac{-g^{\mu\nu} + \frac{q^\mu q^\nu}{m_{Z,\gamma}^2}}{q^2 - m_{Z,\gamma}^2 + im_{Z,\gamma}\Gamma_Z}.$$

For the definitions of the constants used here, the reader is referred to Appendix A.

5.4 Cross Sections

To be able to calculate the differential cross sections, and hence, the total cross section, we need first to obtain the squared matrix element for each Feynman diagram, where use of the trace theorems was made. Later an average over the initial spin polarizations of the electron and the positron pair and the sum over the final spin states of the outgoing particles arising from each initial spin state is carried out. The total cross section as a function of the center of mass energy (see Appendix B) is then calculated.

The calculations were done for the following cases:

$\tan\beta = 05$, $\tan\beta = 15$ and $\tan\beta = 35$ where $M_2 = 150$ or $M_2 = 300$ for each case of $\tan\beta$. All results are given in the following figures.




Figure 5.3: Cross sections for diagram no. 1 in figure 5.1




Figure 5.4: Cross sections for diagram no. 2 in figure 5.1




Figure 5.5: Cross sections for diagram no. 3 in figure 5.1




Figure 5.6: Cross sections for diagram no. 4 in figure 5.1




Figure 5.7: Cross sections for diagram no. 5 in figure 5.1




Figure 5.8: Cross sections for diagram no. 6 in figure 5.1




Figure 5.9: Cross sections for diagram no. 7 in figure 5.1




Figure 5.10: Cross sections for diagram no. 8 in figure 5.1




Figure 5.11: Cross sections for diagram no. 9 in figure 5.1




Figure 5.12: Cross sections for diagram no. 10 in figure 5.1




Figure 5.13: Cross sections for diagram no. 11 in figure 5.1




Figure 5.14: Cross sections for diagram no. 12 in figure 5.1




Figure 5.15: Cross sections for diagram no. 13 in figure 5.1




Figure 5.16: Cross sections for diagram no. 14 in figure 5.1




Figure 5.17: Cross sections for diagram no. 15 in figure 5.2




Figure 5.18: Cross sections for diagram no. 16 in figure 5.2




Figure 5.19: Cross sections for diagram no. 17 in figure 5.2




Figure 5.20: Cross sections for diagram no. 18 in figure 5.2




Figure 5.21: Cross sections for diagram no. 19 in figure 5.2




Figure 5.22: Cross sections for diagram no. 20 in figure 5.2




Figure 5.23: Cross sections for diagram no. 21 in figure 5.2




Figure 5.24: Cross sections for diagram no. 22 in figure 5.2




Figure 5.25: Cross sections for diagram no. 23 in figure 5.2




Figure 5.26: Cross sections for diagram no. 24 in figure 5.2




Figure 5.27: Cross sections for diagram no. 25 in figure 5.2




Figure 5.28: Cross sections for diagram no. 26 in figure 5.2




Figure 5.29: Cross sections for diagram no. 27 in figure 5.2




Figure 5.30: Cross sections for diagram no. 28 in figure 5.2




Figure 5.31: Total cross section for the reaction $e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^0(p5)$

5.5 Conclusion

Results of the previous section are summarized in tables 5.1 and 5.2 for $M_2 = 150$ GeV, and tables 5.3 and 5.4 for $M_2 = 300$ GeV.

From these results, it is clear that the reaction is most probably to proceed through diagram no. 5 through the exchange of a Z boson. The maximum cross section achieved reached the value of 3.4611×10^{-4} [pb] for $\tan\beta = 5$ and $M_2 = 150$ GeV at a center of mass energy, $E_{CM} = 560$ GeV. A maximum of 3.0094×10^{-4} [pb] was obtained for $\tan\beta = 15$ and $M_2 = 150$ GeV at a center of mass energy, $E_{CM} = 560$ GeV and again for the diagram no. 5. For $\tan\beta = 35$ and $M_2 = 150$ GeV at a center of mass energy, $E_{CM} = 580$ GeV, the cross section obtained, reached the value of 2.5145×10^{-4} [pb].

The maximum cross section achieved reached the value of 1.6853×10^{-5} [pb] for $\tan\beta = 5$ and $M_2 = 300$ GeV at a center of mass energy, $E_{CM} = 880$ GeV for the diagram no. 2 which proceeds through the photon propagator, γ . A maximum of 1.0836×10^{-4} [pb] was obtained for $\tan\beta = 15$ and $M_2 = 300$ GeV at a center of mass energy, $E_{CM} = 860$ GeV for the diagram no. 28 which proceeds through the exchange of the scalar neutrino propagator, $\tilde{\nu}_e$. For $\tan\beta = 35$ and $M_2 = 300$ GeV at a center of mass energy, $E_{CM} = 900$ GeV, the cross section obtained, reached the value of 1.8623×10^{-4} [pb] for diagram no. 28.

total cross section achieved the following values,

1. For $\tan\beta = 5$ and $M_2 = 150$, $\sigma_{max} = 7.8383 \times 10^{-4}$ [pb] at $E_{CM} = 720$ GeV.
2. For $\tan\beta = 15$ and $M_2 = 150$, $\sigma_{max} = 5.2422 \times 10^{-4}$ [pb] at $E_{CM} = 820$ GeV.

3. For $\tan\beta = 35$ and $M_2 = 150$, $\sigma_{max} = 4.3095 \times 10^{-4}$ [pb] at $E_{CM} = 800$ GeV.
4. For $\tan\beta = 5$ and $M_2 = 300$, $\sigma_{max} = 3.3550 \times 10^{-4}$ [pb] at $E_{CM} = 1080$ GeV.
5. For $\tan\beta = 15$ and $M_2 = 300$, $\sigma_{max} = 8.8536 \times 10^{-5}$ [pb] at $E_{CM} = 960$ GeV.
6. For $\tan\beta = 35$ and $M_2 = 300$, $\sigma_{max} = 1.3291 \times 10^{-4}$ [pb] at $E_{CM} = 940$ GeV.

Figure No.	$\sigma_{\tan\beta=5}$		$\sigma_{\tan\beta=15}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	1040	1.5779e-06	1140	1.5906e-06	1160	1.6421e-06
2	900	2.0289e-06	880	2.6311e-06	920	2.7928e-06
3	1080	2.2854e-07	1140	2.3085e-07	1120	2.3836e-07
4	860	1.8844e-06	840	2.4976e-06	860	2.6618e-06
5	560	0.00034611	560	0.00030094	580	0.00025145
6	1000	5.3209e-09	1060	7.9594e-09	1060	8.5888e-09
7	1000	2.2084e-10	1060	5.1532e-10	1060	6.4742e-10
8	520	0.000346	560	6.9141e-05	600	3.2496e-05
9	720	1.9348e-05	720	3.2102e-05	740	3.167e-05
10	1480	8.8544e-08	1500	6.5752e-08	1480	5.2112e-08
11	1420	1.2839e-06	1420	9.6548e-07	1460	7.9427e-07
12	780	6.4527e-07	880	2.0701e-07	940	1.1992e-07
13	800	2.6592e-05	800	2.125e-05	780	1.6044e-05
14	1160	1.0001e-08	1160	9.3249e-10	1160	1.4909e-10
15	1200	3.5258e-09	1020	3.4863e-09	1040	3.04e-09
16	1080	7.6575e-08	1040	6.8391e-08	1040	5.8846e-08

Table 5.1: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^0(p5) \text{ for } M_2 = 150 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=5}$		$\sigma_{\tan\beta=15}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
17	760	6.1116e-07	800	1.6734e-07	820	8.3768e-08
18	800	2.6135e-05	780	2.0918e-05	800	1.5747e-05
19	1180	1.3204e-08	1060	3.0367e-08	1040	4.1023e-08
20	1060	1.1262e-07	1060	1.3572e-07	1020	1.3639e-07
21	780	9.6615e-07	800	4.0522e-07	820	2.5265e-07
22	780	3.5148e-05	760	4.295e-05	780	4.0162e-05
23	1440	3.3033e-07	1420	5.7192e-07	1460	7.0432e-07
24	1440	1.8847e-06	1420	1.9127e-06	1460	1.8439e-06
25	780	1.023e-06	880	5.0101e-07	960	3.6067e-07
26	820	3.5797e-05	820	4.3613e-05	820	4.0866e-05
27	1120	4.4032e-07	1040	7.3516e-07	1080	8.5312e-07
28	800	0.00033039	780	0.00023661	800	0.00018492

Table 5.2: Cont. Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^o(p5) \text{ for } M_2 = 150 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=5}$		$\sigma_{\tan\beta=15}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	1460	2.8416e-06	1580	2.6736e-06	1640	2.6225e-06
2	880	1.6853e-05	880	2.2289e-05	860	2.2811e-05
3	1560	4.1064e-07	1560	3.8557e-07	1560	3.7849e-07
4	860	8.5308e-06	860	1.1303e-05	840	1.1522e-05
5	720	6.843e-06	720	4.6316e-05	700	7.9708e-05
6	1280	2.0786e-09	1300	2.293e-09	1360	2.25e-09
7	1280	8.5288e-10	1300	9.6192e-10	1260	9.378e-10
8	980	5.045e-07	980	4.6375e-07	960	4.2704e-07
9	1060	2.0822e-08	1100	2.7708e-08	1120	2.7887e-08
10	1660	3.6361e-07	1560	5.1425e-07	1680	5.5852e-07
11	1520	1.6809e-07	1460	1.5064e-07	1540	1.4015e-07
12	1460	6.7452e-09	1400	1.1721e-08	1400	1.3222e-08
13	1220	4.4132e-06	1280	5.285e-06	1280	5.3928e-06
14	1260	5.8849e-09	1280	8.0866e-10	1280	1.5408e-10
15	1620	9.3888e-09	1220	1.6336e-08	1240	2.0865e-08
16	1200	1.8497e-08	1120	2.1328e-08	1140	2.061e-08

Table 5.3: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^0(p5) \text{ for } M_2 = 300 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=5}$		$\sigma_{\tan\beta=15}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
17	1180	1.1424e-09	1120	2.5658e-09	1100	3.0713e-09
18	1020	1.7106e-06	1060	2.12e-06	1080	2.1515e-06
19	1640	3.0247e-07	1240	4.1924e-07	1200	5.0226e-07
20	1160	1.0304e-08	1140	8.1575e-09	1140	7.0367e-09
21	1160	1.2888e-08	1120	2.5779e-08	1160	2.9906e-08
22	1020	4.1835e-07	1040	3.9874e-07	1060	3.7029e-07
23	1600	1.1700e-05	1660	1.3191e-05	1640	1.3441e-05
24	1540	9.3783e-08	1520	5.7649e-08	1580	4.7951e-08
25	1480	7.6122e-08	1400	1.1802e-07	1480	1.2841e-07
26	1300	1.0781e-06	1300	9.9456e-07	1320	9.3037e-07
27	1040	3.9654e-06	1040	6.3122e-06	1060	6.788e-06
28	880	1.5994e-05	860	0.00010836	900	0.00018623

Table 5.4: Cont. Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow H^-(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^o(p5) \text{ for } M_2 = 300 \text{ GeV}$$

CHAPTER 6

PRODUCTION OF A LIGHT NEUTRAL HIGGS BOSON WITH A CHARGINO AND A NEUTRALINO

6.1 Introduction

In this chapter, the production of a light neutral Higgs boson is considered through the reaction, $e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^-(p5)$, for different topologies and different propagators (see Appendix A). There are a total of 13 feynman diagrams for this reaction (tree level approximation) for which we gave the matrix element corresponding to each diagram. Again, diagrams with the same topology which can be obtained by interchanging the indices were represented once. Our work will proceed as before,

1. Feynman diagrams are given,
2. Diagrams with the same topology are represented once, but has been taken into considerations when calculating the cross section.
3. Matrix elements are written, all the four momenta squares are defined to be mass squared (> 0),
4. Matrix elements are squared,

5. An average over the initial spin polarizations of the electron and positron pair and a sum over the final spin states of the outgoing particles arising from each initial spin state is carried out.
6. Results are represented graphically, and summarized in subsequent tables.

6.2 Feynman Diagrams

The following is the set of Feynman diagrams which were used to calculate the cross section of the associated production of a charged Higgs boson with a chargino and a neutralino. Our momentum notation is: $e^-(p1)$, $e^+(p2)$, $h(p3)$, $\tilde{\chi}_1^+(p4)$ and $\tilde{\chi}_1^-(p5)$.




Figure 6.1: Feynman diagrams for the reaction: $e^-(p1)$, $e^+(p2)$, $h(p3)$ $\tilde{\chi}_1^+(p4)$ and $\tilde{\chi}_1^-(p5)$

6.3 Matrix Elements

The following is the set of matrix elements corresponding to diagrams in figure 6.1 used in our calculations:

$$\mathcal{M}_1 = \bar{v}(p_2)\gamma_\mu u(p_1) \frac{ig e^2}{(p_1 + p_2)^2 + i\epsilon} \bar{u}(p_4) (C_{11}^L P_L + C_{11}^R P_R) \frac{(\not{p}_3 + \not{p}_4 + m_{\tilde{\chi}_1^+})}{(p_3 + p_4)^2 + m_{\tilde{\chi}_1^+}^2 + i\epsilon} \gamma^\mu v(p_5)$$

$$\mathcal{M}_2 = \bar{v}(p_2)\gamma_\mu u(p_1) \frac{ig e^2}{(p_1 + p_2)^2 + i\epsilon} \bar{u}(p_5)\gamma^\mu \frac{(\not{p}_3 + \not{p}_5 + m_{\tilde{\chi}_1^-})}{(p_3 + p_5)^2 + m_{\tilde{\chi}_1^-}^2 + i\epsilon} (C_{11}^L P_L + C_{11}^R P_R) v(p_4)$$

$$\begin{aligned} \mathcal{M}_3 = & \bar{v}(p_2)\gamma_\sigma(g_V - \gamma_5)u(p_1) \frac{ig^3 M_Z \sin(\beta - \alpha)}{4 \cos^3 \theta_w} \bar{u}(p_5)\gamma^\mu (O_{11}^L P_L + O_{11}^R P_R) v(p_4) \\ & \times \frac{(g_{\mu\nu} - q_\mu q_\nu / m_Z^2)(g^{\nu\sigma} - k^\nu k^\sigma / M_Z^2)}{((p_1 + p_2 - p_5)^2 - M_Z^2 + i\epsilon)} \end{aligned}$$

$$\begin{aligned} \mathcal{M}_4 = & \bar{v}(p_2)\gamma^\nu(g_V - \gamma_5)u(p_1) \frac{ig^3 \cos(\alpha - \beta)}{8 \cos^2 \theta_w} \bar{u}(p_5) (C_{11}^{A,L} P_L + C_{11}^{A,R} P_R) v(p_4) \\ & \times \frac{(q_\mu - h_\mu)}{q^2 - M_{H_3}^2 + i\epsilon} \frac{(g^{\mu\nu} - k^\mu k^\nu / M_Z^2)}{((p_1 + p_2)^2 - M_Z^2 + i\epsilon)} \end{aligned}$$

$$\begin{aligned} \mathcal{M}_5 = & \bar{v}(p_2)\gamma^\nu(g_V - \gamma_5)u(p_1) \frac{-ig^3}{4 \cos^2 \theta_w ((p_1 + p_2)^2 - M_Z^2 + i\epsilon)} \bar{u}(p_4) (C_{11}^L P_L + C_{11}^R P_R) \\ & \times \frac{(\not{p}_3 + \not{p}_4 + m_{\tilde{\chi}_1^+})}{(p_3 + p_4)^2 + m_{\tilde{\chi}_1^+}^2 + i\epsilon} \gamma^\mu (O_{11}^L P_L + O_{11}^R P_R) v(p_5) \left(g_{\mu\nu} - \frac{k_\mu k_\nu}{M_Z^2} \right) \end{aligned}$$

$$\begin{aligned}\mathcal{M}_6 = & \bar{v}(p_2)\gamma^\nu(g_V - \gamma_5)u(p_1)\frac{-ig^3}{4\cos^2\theta_w((p_1+p_2)^2 - M_Z^2 + i\epsilon)}\bar{u}(p_5)\gamma^\mu(O_{11}^L P_L + O_{11}^R P_R) \\ & \times \frac{-(\not{p}_3' + \not{p}_5' + m_{\tilde{\chi}_1^+})}{(p_3 + p_5)^2 + m_{\tilde{\chi}_1^+}^2 + i\epsilon}(C_{11}^L P_L + C_{11}^R P_R)v(p_4)\left(g_{\mu\nu} - \frac{k_\mu k_\nu}{M_Z^2}\right)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_7 = & \bar{v}(p_2)\gamma^\nu(g_V - \gamma_5)u(p_1)\frac{-ig^3}{4\cos^2\theta_w((p_1+p_2)^2 - M_Z^2 + i\epsilon)}\bar{u}(p_4)(C_{12}^L P_L + C_{12}^R P_R) \\ & \times \frac{\not{p}_3' + \not{p}_4' + m_{\tilde{\chi}_2^+}}{(p_3 + p_4)^2 + m_{\tilde{\chi}_2^+}^2 + i\epsilon}\gamma^\mu(O_{21}^L P_L + O_{21}^R P_R)\left(g_{\mu\nu} - \frac{k_\mu k_\nu}{M_Z^2}\right)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_8 = & \bar{v}(p_2)\gamma^\nu(g_V - \gamma_5)u(p_1)\frac{-ig^3}{4\cos^2\theta_w((p_1+p_2)^2 - M_Z^2 + i\epsilon)}\bar{u}(p_5)\gamma^\mu(O_{12}^L P_L + O_{12}^R P_R) \\ & \times \frac{\not{p}_3' + \not{p}_5' + m_{\tilde{\chi}_2^+}}{(p_3 + p_5)^2 + m_{\tilde{\chi}_2^+}^2 + i\epsilon}(C_{21}^L P_L + C_{21}^R P_R)v(p_4)\left(g_{\mu\nu} - \frac{k_\mu k_\nu}{M_Z^2}\right)\end{aligned}$$

$$\mathcal{M}_9 = \frac{ig^3|V_{11}|^2}{(p_1 - p_5)^2 - m_\nu^2}\bar{v}(p_2)P_L\frac{\not{p}_3' + \not{p}_4' + m_{\tilde{\chi}_1^+}}{((p_3 + p_4)^2 - m_{\tilde{\chi}_1^+}^2 + i\epsilon)}(C_{11}^L P_L + C_{11}^R P_R)u(p_5)\bar{v}(p_4)P_R\bar{u}(p_1)$$

$$\mathcal{M}_{10} = \frac{ig^3|V_{11}||V_{21}|}{(p_1 - p_5)^2 - m_\nu^2}\bar{v}(p_2)P_L\frac{\not{p}_3' + \not{p}_4' + m_{\tilde{\chi}_1^+}}{((p_3 + p_4)^2 - m_{\tilde{\chi}_1^+}^2 + i\epsilon)}(C_{21}^L P_L + C_{21}^R P_R)u(p_5)\bar{v}(p_4)P_R\bar{u}(p_1)$$

$$\mathcal{M}_{11} = \frac{ig^3 M_W \sin(\alpha + \beta)|V_{11}|^2}{2\cos^2\theta_w((p_1 - p_5)^2 - m_\nu^2)((p_1 - p_4)^2 - m_\nu^2)}\bar{v}(p_2)P_L u(p_5)\bar{v}(p_4)P_R\bar{u}(p_1)$$

$$\mathcal{M}_{12} = \frac{ig^3|V_{11}||V_{21}|}{((p_1 - p_4)^2 - m_\nu^2)}\bar{v}(p_2)P_L u(p_5)\bar{v}(p_4)(C_{21}^L P_L + C_{21}^R P_R)\frac{-(\not{p}_3' + \not{p}_5') + m_{\tilde{\chi}_2^+}}{((p_3 + p_5)^2 - m_{\tilde{\chi}_2^+}^2 + i\epsilon)}P_R\bar{u}(p_1)$$

$$\mathcal{M}_{13} = \frac{ig^3 |V_{11}|^2}{((p_1 - p_4)^2 - m_\nu^2)} \bar{v}(p_2) P_L u(p_5) \bar{v}(p_4) (C_{11}^L P_L + C_{11}^R P_R) \frac{-(\not{p}_3 + \not{p}_5) + m_{\tilde{\chi}_1^+}}{((p_3 + p_5)^2 - m_{\tilde{\chi}_1^+}^2 + i\epsilon)} P_R \bar{u}(p_1)$$

For the definitions of the constants used here, the reader is referred to Appendix A.

6.4 Cross Sections

As before, to calculate the differential cross sections, and hence, the total cross section, we need first to obtain the squared matrix element for each Feynman diagram, where use of the trace theorems was made. Later an average over the initial spin polarizations of the electron and the positron pair and the sum over the final spin states of the outgoing particles arising from each initial spin state is carried out. The total cross section as a function of the center of the mass energy (see Appendix B) is then calculated.

Calculations were done with the following set of parameters:

$\tan\beta = 10$, and $\tan\beta = 15$ where $M_2 = 150$ or $M_2 = 300$.

All results are given in the following figures.




Figure 6.2: Cross sections for diagram no. 1 in figure 6.1




Figure 6.3: Cross sections for diagram no. 2 in figure 6.1




Figure 6.4: Cross sections for diagram no. 3 in figure 6.1




Figure 6.5: Cross sections for diagram no. 4 in figure 6.1




Figure 6.6: Cross sections for diagram no. 5 in figure 6.1




Figure 6.7: Cross sections for diagram no. 6 in figure 6.1




Figure 6.8: Cross sections for diagram no. 7 in figure 6.1




Figure 6.9: Cross sections for diagram no. 8 in figure 6.1




Figure 6.10: Cross sections for diagram no. 09 in figure 6.1




Figure 6.11: Cross sections for diagram no. 10 in figure 6.1




Figure 6.12: Cross sections for diagram no. 11 in figure 6.1




Figure 6.13: Cross sections for diagram no. 12 in figure 6.1





Figure 6.14: Cross sections for diagram no. 13 in figure 6.1




Figure 6.15: Total cross sections for the reaction $e^- e^+ \rightarrow h \tilde{\chi}_1^+ \tilde{\chi}_1^-$

6.5 Conclusion

Results of the previous section are summarized in tables 6.1 and 6.2 for $M_2 = 150$ GeV and $M_2 = 300$ GeV respectively.

From these results, it was found that the maximum cross section obtained is 2.0371×10^{-3} [pb] at $\tan\beta = 10$ and $M_2 = 150$ GeV for diagram number 7, in which, the reaction proceeds through the Z boson propagator. It takes place at a center of mass energy, $E_{CM} = 560$ GeV. For $\tan\beta = 15$, the maximum cross section is 1.8340×10^{-3} [pb] where $M_2 = 150$ GeV for the diagram no. 8, in which, the reaction proceeds, again, through the Z boson propagator. It occurs at a center of mass energy, $E_{CM} = 560$ GeV.

The maximum cross section obtained by this reaction for $\tan\beta = 10$ and $M_2 = 300$ GeV is 1.8605×10^{-3} [pb] at a center of mass energy, $E_{CM} = 580$ GeV for the feynman diagram no. 7 which proceeds through the Z boson propagator. While the maximum cross section takes the value 5.4781×10^{-5} [pb] for $\tan\beta = 15$ and $M_2 = 300$ GeV and at a center of mass energy, $E_{CM} = 960$ GeV. It occurs at diagram no. 1, which proceeds through the photon propagator.

Total cross section of this reaction is 3.0430×10^{-3} [pb] for $\tan\beta = 10$ and $M_2 = 150$ GeV at a center of mass energy, $E_{CM} = 780$ GeV, and is 1.4490×10^{-4} [pb] for $\tan\beta = 15$ and $M_2 = 150$ GeV at a center of mass energy, $E_{CM} = 920$ GeV.

Total cross section also assumes the value 2.7681×10^{-3} [pb] for $\tan\beta = 15$ and $M_2 = 300$ GeV at a center of mass energy, $E_{CM} = 840$ GeV, and 1.0993×10^{-4} [pb] for $\tan\beta = 15$ and $M_2 = 300$ GeV at a center of mass energy, $E_{CM} = 920$ GeV.

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=15}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	660	2.8134e-05	960	6.5301e-05
2	660	2.819e-05	960	6.5243e-05
3	600	1.1079e-05	1240	1.1034e-06
4	1000	3.7658e-07	1720	1.8818e-07
5	660	2.8988e-05	960	3.4915e-05
6	660	2.9043e-05	960	3.4883e-05
7	560	0.0020371	720	1.1311e-05
8	560	0.0020166	560	0.001834
9	1180	3.7073e-05	1160	2.8931e-05
10	860	0.0017111	800	0.0014741
11	1100	5.2035e-07	1120	5.8349e-07
12	1140	3.6974e-05	1100	2.885e-05
13	780	0.0016977	860	0.0014679
total	780	0.003043	920	0.0001493

Table 6.1: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^-(p5) \text{ for } M_2 = 150 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=15}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	680	2.1275e-05	960	5.4781e-05
2	660	2.1232e-05	960	5.4754e-05
3	600	1.0909e-05	1340	1.1712e-06
4	1060	2.1442e-07	1820	1.0861e-07
5	680	2.1987e-05	960	2.9262e-05
6	660	2.1988e-05	960	2.9257e-05
7	580	0.0018605	780	2.6224e-06
8	720	1.118e-05	800	2.6398e-06
9	1280	4.6013e-05	1260	3.9595e-05
10	840	2.8019e-05	1100	8.4166e-06
11	1280	1.6293e-07	1340	1.8258e-07
12	1280	4.5969e-05	1260	3.962e-05
13	920	2.8152e-05	1160	8.646e-06
total	840	0.0027681	920	0.00010993

Table 6.2: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^+(p4)\tilde{\chi}_1^-(p5) \text{ for } M_2 = 300 \text{ GeV}$$

CHAPTER 7

PRODUCTION OF A LIGHT NEUTRAL HIGGS BOSON WITH A PAIR OF NEUTRALINOS

7.1 Introduction

In this chapter, the production of a light neutral Higgs boson is considered through the reaction, $e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^o(p4)\tilde{\chi}_1^o(p5)$, for different topologies and different propagators (see Appendix A). There are a total of 24 Feynman diagrams for this reaction (tree level approximation) for which we gave the matrix element corresponding to each diagram. Again, diagrams with the same topology which can be obtained by interchanging the indices were represented once. Our work will proceed as before,

1. Feynman diagrams are given,
2. Diagrams with the same topology are represented once, but has been taken into considerations when calculating the cross section.
3. Matrix elements are written, all the four momenta squares are defined to be mass squared (> 0),
4. Matrix elements are squared,

5. An average over the initial spin polarizations of the electron and positron pair and a sum over the final spin states of the outgoing particles arising from each initial spin state is carried out.
6. Results are represented graphically, and summarized in subsequent tables.

7.2 Feynman Diagrams

The following is the set of Feynman diagrams which were used to calculate the cross section of the associated production of a charged Higgs boson with a pair of neutralinos. Our momentum notation is: $e^-(p1)$, $e^+(p2)$, $h(p3)$ $\tilde{\chi}_1^o(p4)$ and $\tilde{\chi}_1^o(p5)$.




Figure 7.1: Feynman diagrams for the reaction: $e^-(p_1)e^+(p_2) \rightarrow h(p_3)\tilde{\chi}_1^o(p_4)\tilde{\chi}_1^o(p_5)$




Figure 7.2: Cont. Feynman diagrams for the reaction: $e^-(p_1)e^+(p_2) \rightarrow h(p_3)\tilde{\chi}_1^o(p_4)\tilde{\chi}_1^o(p_5)$

7.3 Matrix Elements

The following is the set of matrix elements corresponding to diagrams in figures 7.1 and 7.2 used in our calculations:

$$\begin{aligned}\mathcal{M}_1 = & -i\bar{v}(p_2)\gamma^\mu(B^L P_L + B^R P_L)u(p_1)\frac{g^{\mu\nu} - k_\mu k_\nu/M_Z^2}{(p_1 + p_2)^2 - M_Z^2 + i\epsilon} \\ & J_1 g^{\nu\rho}\frac{g^{\rho\sigma} - k_\rho k_\sigma/M_Z^2}{(p_4 + p_5)^2 - M_Z^2 + i\epsilon} \times \bar{u}(p_5)\gamma_\sigma(S_{11} + S_{11}'\gamma_5)v(p_4)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_2 = & i\bar{v}(p_2)\gamma^\mu(B^L P_L + B^R P_L)u(p_1)\frac{g^{\mu\nu} - k_\mu k_\nu/M_Z^2}{(p_1 + p_2)^2 - M_Z^2 + i\epsilon} \times \\ & G_1(p_4 + p_5 - p_3)^\nu\frac{1}{(p_4 + p_5)^2 - m_{H_3}^2}\bar{u}(p_4)(R_{311} + R_{311}'\gamma_5)v(p_5)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_{3,4,5,6} = & \bar{v}(p_2)\gamma^\mu(B^L P_L + B^R P_L)u(p_1)\frac{g^{\mu\nu} - p_1^\mu p_2^\nu/M_Z^2}{(p_1 + p_2)^2 - M_Z^2 + i\epsilon}\bar{u}(p_5)(R_{111} + R_{111}'\gamma_5) \\ & \times \frac{\not{p}_3 + \not{p}_5 + m_{\tilde{\chi}_1^0}}{(p_3 + p_5)^2 - m_{\tilde{\chi}_1^0}^2 + i\epsilon}\gamma_\nu(S_{11} + S_{11}'\gamma_5)v(p_4)\end{aligned}$$

where k assumes the values, 1, 2, 3, 4.

$$\begin{aligned}\mathcal{M}_{7,8,9,10} = & -i\bar{v}(p_2)(N_{L1} + N_{L1}'\gamma_5)\frac{\not{p}_3 + \not{p}_4 + m_{\tilde{\chi}_k^0}}{(p_3 + p_4)^2 - m_{\tilde{\chi}_k^0}^2 + i\epsilon}(R_{111} + R_{111}'\gamma_5)u(p_4) \\ & \times \frac{\not{p}_1 - \not{p}_5 + m_{\tilde{e}}}{(p_1 - p_5)^2 - m_{\tilde{e}}^2 + i\epsilon}\bar{u}(p_5)(N_{L1} + N_{L1}'\gamma_5)u(p_1)\end{aligned}$$

again, here k assumes the values, 1, 2, 3, 4.

$$\begin{aligned}\mathcal{M}_{11} = & -\bar{v}(p_2)(N_{L1} + N_{L1}'\gamma_5)u(p_5)\frac{\not{p}_2 - \not{p}_5 + m_{\tilde{e}}}{(p_2 - p_5)^2 - m_{\tilde{e}}^2 + i\epsilon}\psi_k\frac{\not{p}_1 - \not{p}_4 + m_{\tilde{e}}}{(p_1 - p_4)^2 - m_{\tilde{e}}^2 + i\epsilon} \\ & \times \bar{u}(p_4)(N_{L1} + N_{L1}'\gamma_5)u(p_1)\end{aligned}$$

$$\text{Here, } \psi = \frac{igM_Z \cos(\alpha+\beta)}{\cos\theta_w}\left(\frac{1}{2} + \sin^2\theta_w\right).$$

$$\begin{aligned}\mathcal{M}_{12,13,14,15} = & -i\bar{v}(p_2)(N_{L1} + N_{L1}'\gamma_5)u(p_4)\frac{\not{p}_2 - \not{p}_4 + m_{\tilde{e}}}{(p_2 - p_4)^2 - m_{\tilde{e}}^2 + i\epsilon}\bar{u}(p_5)(R_{111} + R_{111}'\gamma_5) \\ & \times \frac{\not{p}_3 + \not{p}_5 + m_{\tilde{\chi}_k^0}}{(p_3 + p_5)^2 - m_{\tilde{\chi}_k^0}^2 + i\epsilon}(N_{L1} + N_{L1}'\gamma_5)u(p_1)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_{16,17,18,19} = & -i\bar{v}(p_2)(N_{R1} + N_{R1}'\gamma_5)\frac{\not{p}_1 - \not{p}_5 + m_{\tilde{\chi}_k^0}}{(p_1 - p_5)^2 - m_{\tilde{\chi}_k^0}^2 + i\epsilon}(R_{111} + R_{111}'\gamma_5)u(p_4) \\ & \times \frac{\not{p}_1 - \not{p}_5 + m_{\tilde{e}}}{(p_1 - p_5)^2 - m_{\tilde{e}}^2 + i\epsilon}\bar{u}(p_5)(N_{R1} + N_{R1}'\gamma_5)u(p_1)\end{aligned}$$

As above, k takes the vlaues, 1, 2, 3, 4.

$$\begin{aligned}\mathcal{M}_{20} = & -\bar{v}(p_2)(N_{R1} + N_{R1}'\gamma_5)u(p_4)\frac{\not{p}_2 - \not{p}_4 + m_{\tilde{e}}}{(p_2 - p_4)^2 - m_{\tilde{e}}^2 + i\epsilon}\psi_k\frac{\not{p}_1 - \not{p}_5 + m_{\tilde{e}}}{(p_1 - p_5)^2 - m_{\tilde{e}}^2 + i\epsilon} \\ & \times \bar{u}(p_5)(N_{R1} + N_{R1}'\gamma_5)u(p_1)\end{aligned}$$

$$\text{and also, } \psi = \frac{igM_Z \cos(\alpha+\beta)}{\cos\theta_w}\left(\frac{1}{2} + \sin^2\theta_w\right).$$

$$\begin{aligned}\mathcal{M}_{21,22,23,24} = & \bar{v}(p_2)(N_{R1} + N_{R1}'\gamma_5)u(p_5)\frac{\not{p}_2 - \not{p}_5 + m_{\tilde{e}}}{(p_2 - p_5)^2 - m_{\tilde{e}}^2 + i\epsilon}\bar{u}(p_4)(R_{111} + R_{111}'\gamma_5) \\ & \times (R_{111} + R_{111}'\gamma_5)\frac{\not{p}_3 + \not{p}_4 + m_{\tilde{\chi}_k^0}}{(p_3 + p_4)^2 - m_{\tilde{\chi}_k^0}^2 + i\epsilon}(N_{R1} + N_{R1}'\gamma_5)u(p_1)\end{aligned}$$

and $k = 1, 2, 3, 4$.

For the definitions of the constants used here, the reader is referred to Appendix A.

7.4 Cross Sections

As before, to calculate the differential cross sections, and hence, the total cross section, we need first to obtain the squared matrix element for each Feynman diagram, where use of the trace theorems was made. Later an average over the initial spin polarizations of the electron and the positron pair and the sum over the final spin states of the outgoing particles arising from each initial spin state is carried out. The total cross section as a function of the center of the mass energy (see Appendix B) is then calculated.

Calculations were done with the following set of parameters:

$\tan\beta = 10$, and $\tan\beta = 15$ where $M_2 = 150$ or $M_2 = 300$.

All results are given in the following figures.




Figure 7.3: Cross sections for diagram no. 1 in figure 7.1




Figure 7.4: Cross sections for diagram no. 2 in figure 7.1




Figure 7.5: Cross sections for diagram no. 3 in figure 7.1




Figure 7.6: Cross sections for diagram no. 4 in figure 7.1





Figure 7.7: Cross sections for diagram no. 5 in figure 7.1




Figure 7.8: Cross sections for diagram no. 6 in figure 7.1




Figure 7.9: Cross sections for diagram no. 7 in figure 7.1




Figure 7.10: Cross sections for diagram no. 8 in figure 7.1





Figure 7.11: Cross sections for diagram no. 9 in figure 7.1




Figure 7.12: Cross sections for diagram no. 10 in figure 7.1





Figure 7.13: Cross sections for diagram no. 11 in figure 7.1




Figure 7.14: Cross sections for diagram no. 12 in figure 7.1





Figure 7.15: Cross sections for diagram no. 13 in figure 7.1





Figure 7.16: Cross sections for diagram no. 14 in figure 7.1




Figure 7.17: Cross sections for diagram no. 15 in figure 7.2




Figure 7.18: Cross sections for diagram no. 16 in figure 7.2





Figure 7.19: Cross sections for diagram no. 17 in figure 7.2




Figure 7.20: Cross sections for diagram no. 18 in figure 7.2




Figure 7.21: Cross sections for diagram no. 19 in figure 7.2




Figure 7.22: Cross sections for diagram no. 20 in figure 7.2





Figure 7.23: Cross sections for diagram no. 21 in figure 7.2




Figure 7.24: Cross sections for diagram no. 22 in figure 7.2





Figure 7.25: Cross sections for diagram no. 23 in figure 7.2




Figure 7.26: Cross sections for diagram no. 24 in figure 7.2




Figure 7.27: Total cross sections for the reaction $e^- e^+ \rightarrow h \tilde{\chi}_1^o \tilde{\chi}_1^o$

7.5 Conclusion

Results of the previous section are summarized in tables 7.1 and 7.2 for $M_2 = 150$ GeV and in tables 7.3 and 7.4 for $M_2 = 300$ GeV respectively. From these results, it is clear that the reaction is most probably to proceed through Feynman diagram no. 5. The cross section of this reaction achieved a value of 1.5336×10^{-4} [pb] at $\tan\beta = 10$ and for $M_2 = 150$ GeV. And the cross section takes the value of 2.3852×10^{-4} [pb] at $\tan\beta = 35$ and for $M_2 = 150$ GeV. for $M_2 = 300$ GeV, the value of the cross section achieved 3.0865×10^{-4} [pb] at $\tan\beta = 10$, and the value 4.0505×10^{-4} [pb] at $\tan\beta = 35$. The total cross section of this reaction reaches its maximum value, 8.3901×10^{-4} [pb], at $\tan\beta = 35$ and for $M_2 = 150$ GeV. Interference effects which are not studied here, should be taken into accounts when dealing with the total cross section. for example the interference terms emerging from diagrams 20 & 5 ($M_2 = 150\text{GeV}$) would decrease the value of the total cross section effectively.

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	1000	4.2976e-07	1060	4.8279e-07
2	920	3.0571e-07	980	4.8221e-08
3	820	8.7088e-08	840	4.3811e-08
4	760	8.4636e-11	800	1.1105e-10
5	520	0.00015336	540	0.00023852
6	720	9.8329e-05	740	0.00010254
7	1300	1.7996e-07	1340	5.4578e-08
8	1260	4.3651e-08	1340	2.7638e-08
9	800	3.3079e-07	800	5.2466e-07
10	780	7.8608e-05	780	5.0584e-05
11	1260	8.1842e-10	1280	6.0505e-10
12	1300	1.7990e-07	1340	5.45e-08
13	1240	4.3619e-08	1340	2.7632e-08
14	780	3.3081e-07	800	5.2398e-07
15	800	7.8692e-05	800	5.0708e-05
16	1280	1.176e-06	1360	7.3614e-07

Table 7.1: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^o(p4)\tilde{\chi}_1^o(p5) \text{ for } M_2 = 150 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
17	1280	7.8242e-08	1320	6.4104e-08
18	780	6.9236e-07	800	1.581e-06
19	780	0.00013973	800	0.0001293
20	1260	3.9622e-09	1280	6.0674e-09
21	1340	1.1762e-06	1340	7.3803e-07
22	1240	7.8012e-08	1340	6.4162e-08
23	780	6.9132e-07	780	1.576e-06
24	780	0.00013979	800	0.00012911

Table 7.2: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^o(p4)\tilde{\chi}_1^o(p5) \text{ for } M_2 = 150 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	1080	5.3869e-08	1120	5.5643e-08
2	1040	3.0443e-08	1020	4.506e-09
3	860	1.4074e-09	880	6.0213e-10
4	680	2.7783e-07	680	1.6657e-07
5	540	0.00011284	540	0.00017088
6	800	3.8825e-06	800	5.3689e-06
7	1380	4.5216e-08	1380	2.3421e-08
8	880	7.155e-06	940	3.9801e-06
9	780	3.5995e-07	800	8.0505e-07
10	840	0.00030865	840	0.00040393
11	1260	1.4936e-09	1300	2.0679e-09
12	1340	4.5271e-08	1360	2.3366e-08
13	880	7.1686e-06	940	3.9779e-06
14	800	3.5916e-07	800	8.0607e-07
15	860	0.00030897	860	0.00040505
16	1300	1.2266e-06	1340	5.621e-07

Table 7.3: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^o(p4)\tilde{\chi}_1^o(p5) \text{ for } M_2 = 300 \text{ GeV}$$

Figure No.	$\sigma_{tan,\beta=10}$		$\sigma_{tan,\beta=35}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
17	940	3.0376e-06	940	1.3584e-06
18	800	3.7196e-06	800	7.8338e-06
19	820	6.2326e-05	840	6.9613e-05
20	1260	3.0118e-08	1300	3.6928e-08
21	1360	1.2275e-06	1340	5.627e-07
22	880	3.0427e-06	940	1.3604e-06
23	780	3.7302e-06	800	7.8293e-06
24	840	6.2285e-05	860	6.9524e-05

Table 7.4: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^o(p4)\tilde{\chi}_1^o(p5) \text{ for } M_2 = 300 \text{ GeV}$$

CHAPTER 8

PRODUCTION OF A LIGHT NEUTRAL HIGGS BOSON WITH A PAIR OF CHARGED HIGGS BOSONS

8.1 Introduction

In this chapter, the production of a light neutral Higgs boson is considered through the reaction, $e^-(p1)e^+(p2) \rightarrow h(p3)H^+(p4)H^-(p5)$, for different topologies and different propagators (see Appendix A). There are a total of 5 Feynman diagrams for this reaction (tree level approximation) for which we gave the matrix element corresponding to each diagram. Again, diagrams with the same topology which can be obtained by interchanging the indices were represented once. Our work will proceed as usual,

1. Feynman diagrams are given,
2. Diagrams with the same topology are represented once, but has been taken into considerations when calculating the cross section.
3. Matrix elements are written, all the four momenta squares are defined to be mass squared (> 0),
4. Matrix elements are squared,

5. An average over the initial spin polarizations of the electron and positron pair and a sum over the final spin states of the outgoing particles arising from each initial spin state is carried out.
6. Results are represented graphically, and summarized in subsequent tables.

8.2 Feynman Diagrams

The following is the set of Feynman diagrams which were used to calculate the cross section of the associated production of a charged Higgs boson with a pair of neutralinos. Our momentum notation is: $e^-(p1)$, $e^+(p2)$, $h(p3)$, $H^+(p4)$, and $H^-(p5)$.




Figure 8.1: Feynman diagrams for the reaction: $e^-(p1)e^+(p2) \rightarrow h(p3)\tilde{\chi}_1^o(p4)\tilde{\chi}_1^o(p5)$

8.3 Matrix Elements

The following is the set of matrix elements corresponding to diagrams in figures 8.1 used in our calculations:

$$\begin{aligned}\mathcal{M}_1 = & \bar{v}(\vec{p}_2) A^\mu \gamma_\mu \frac{g^{\mu\nu}}{(p_1 + p_2)^2} e(p_1 + p_2)^\mu \frac{1}{(p_4 + p_5)^2 - m_{H_3}^2} \\ & \times g \left[M_w \cos(\beta - \alpha) - \frac{M_Z}{2 \cos \theta_w} \cos 2\beta \cos(\beta + \alpha) \right] u(p_1)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_2 = & \bar{v}(\vec{p}_2) A^\mu \gamma_\mu \frac{g^{\mu\nu}}{(p_1 + p_2)^2} e(p_1 + p_2)^\mu \frac{1}{(p_3 + p_5)^2 - m_{H_3}^2} \\ & \times g \left[M_w \cos(\beta - \alpha) - \frac{M_Z}{2 \cos \theta_w} \cos 2\beta \cos(\beta + \alpha) \right] u(p_1)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_3 = & \bar{v}(\vec{p}_2) \gamma_\mu (B^L P_L + B^R P_R)^\mu \frac{g^{\mu\nu} - k_\mu k_\nu / M_Z}{(p_1 + p_2)^2 - M_Z^2 + i\epsilon} J_1 g^{\nu\rho} \\ & \times \frac{g^{\rho\sigma} - k_\rho k_\sigma / M_Z}{(p_3 + p_4)^2 - M_Z^2 + i\epsilon} \frac{g \cos 2\theta_w}{2 \cos \theta_w} (p_3 + p_4)^\sigma u(p_1)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_4 = & -\bar{v}(\vec{p}_2) \gamma_\mu (B^L P_L + B^R P_R)^\mu \frac{g^{\mu\nu} - k_\mu k_\nu / M_Z}{(p_1 + p_2)^2 - M_Z^2 + i\epsilon} \frac{g \cos 2\theta_w}{2 \cos \theta_w} (p_3 + p_4 + p_5)^\nu \\ & \times \frac{1}{(p_4 + p_5)^2 - m_{H_3}^2} g \left[M_w \cos(\beta - \alpha) - \frac{M_Z}{2 \cos \theta_w} \cos 2\beta \cos(\beta + \alpha) \right] u(p_1)\end{aligned}$$

$$\begin{aligned}\mathcal{M}_5 = & -\bar{v}(\vec{p}_2) \gamma_\mu (B^L P_L + B^R P_R)^\mu \frac{g^{\mu\nu} - k_\mu k_\nu / M_Z}{(p_1 + p_2)^2 - M_Z^2 + i\epsilon} \frac{g \cos 2\theta_w}{2 \cos \theta_w} (p_3 + p_4 + p_5)^\nu \\ & \times \frac{1}{(p_3 + p_5)^2 - m_{H_3}^2} g \left[M_w \cos(\beta - \alpha) - \frac{M_Z}{2 \cos \theta_w} \cos 2\beta \cos(\beta + \alpha) \right] u(p_1)\end{aligned}$$

For the definitions of the constants used here, the reader is referred to Appendix A.

8.4 Cross Sections

As before, to calculate the differential cross sections, and hence, the total cross section, we need first to obtain the squared matrix element for each Feynman diagram, where use of the trace theorems was made. Later an average over the initial spin polarizations of the electron and the positron pair and the sum over the final spin states of the outgoing particles arising from each initial spin state is carried out. The total cross section as a function of the center of the mass energy (see Appendix B) is then calculated.

Calculations were done with the following set of parameters:

$\tan\beta = 10$, and $\tan\beta = 15$ where $M_2 = 150$ or $M_2 = 300$.

All results are given in the following figures.




Figure 8.2: Cross sections for diagram no. 1 in figure 8.1




Figure 8.3: Cross sections for diagram no. 2 in figure 8.1




Figure 8.4: Cross sections for diagram no. 3 in figure 8.1




Figure 8.5: Cross sections for diagram no. 4 in figure 8.1




Figure 8.6: Cross sections for diagram no. 5 in figure 8.1




Figure 8.7: Total cross section for the reaction $e^-(p_1)e^+(p_2) \rightarrow h(p_3)H^+(p_4)H^-(p_5)$

8.5 Conclusion

Results of the previous section are summarized in tables 8.1 for $M_2 = 150$ GeV and in tables 8.2 for $M_2 = 300$ GeV respectively. From these results, it is clear that the reaction is most probably to proceed through diagram no. 3 at $\tan\beta = 10$ and $\tan\beta = 15$ where $M_2 = 150$ GeV. at these values the maximum cross section achieved is 5.7627×10^{-8} [pb] and 5.8264×10^{-8} [pb] respectively.

The total cross section achieved by this reaction is 1.9263×10^{-7} [pb] at $E_{CM} = 820$ GeV for $\tan\beta = 10$ and $M_2 = 150$ GeV, and is 5.9956×10^{-8} [pb] at $E_{CM} = 840$ GeV for $\tan\beta = 15$ and $M_2 = 300$ GeV the maximum cross section at $M_2 = 300$ GeV for both values of $\tan\beta$, that is 10 & 15, takes the values 5.8647×10^{-8} at $E_{CM} = 800$ GeV, and 5.8883×10^{-8} at $E_{CM} = 820$ GeV.

The total cross section achieved by this reaction is 2.0128×10^{-7} [pb] at $E_{CM} = 860$ GeV for $\tan\beta = 10$ and $M_2 = 150$ GeV, and is 6.6248×10^{-8} [pb] at $E_{CM} = 820$ GeV for $\tan\beta = 15$ and $M_2 = 300$ GeV

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=15}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	940	2.6354e-08	960	2.9167e-11
2	940	2.6350e-08	960	2.9131e-11
3	800	5.7627e-08	800	5.8246e-08
4	940	3.8517e-09	960	4.2596e-12
5	940	3.8512e-09	940	4.2561e-12
total	820	1.9263e-07	840	5.9956e-08

Table 8.1: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)H^+(p4)H^-(p5) \text{ for } M_2 = 150 \text{ GeV}$$

Figure No.	$\sigma_{\tan\beta=10}$		$\sigma_{\tan\beta=15}$	
	E_{CM}	σ (Pb)	E_{CM}	σ (Pb)
1	940	2.8116e-08	960	3.9597e-10
2	940	2.8066e-08	940	3.9717e-10
3	800	5.8047e-08	820	5.8883e-08
4	900	4.1112e-09	960	5.7828e-11
5	940	4.102e-09	940	5.8048e-11
total	860	2.0128e-07	820	6.6248e-08

Table 8.2: Summary of the results obtained for the reaction,

$$e^-(p1)e^+(p2) \rightarrow h(p3)H^+(p4)H^-(p5) \text{ for } M_2 = 300 \text{ GeV}$$

APPENDICES

introduction: In this appendix, we summarize the Feynman rules in a convenient form, required to construct the covariant amplitudes in this study. The given rules are computed by using the unitary (physical) gauge [33], [34], [35]. This gauge is the preferred choice in the tree approximation level. For Higher order corrections, we must include the ghost loops, and in this case, it is much more efficient and easier to use the so called 't Hooft-Feynman gauge [36], [37] than the unitary gauge.

A. Appendix A: Feynman Rules

The Feynman diagrams at the tree level for two particles in the initial state and three particles at the final state are classified physically into three different topologies according to the interaction of the two initial particles (or annihilation of particle-antiparticle pair) through no propagators, one propagator, and two propagators.

External Lines (Incoming and Outgoing Particles)

Fermions

	Incoming	Outgoing
Particle	$u(p, s)$	$\bar{u}(p, s)$
AntiParticle	$\bar{v}(p, s)$	$v(p, s)$

Vector Bosons

Incoming	Outgoing
$\varepsilon_\sigma(p, s)$	$\varepsilon_\sigma^*(p, s)$

Scalar (Higgs) Bosons

Scalar bosons as external lines take the value **1** in general.

Internal Lines (Propagators)

fermions

$$i \frac{\not{p} + m}{p^2 - m^2 + i\epsilon}$$

Massive Vector Bosons

$$-i \frac{g^{\mu\nu} - k^\mu k^\nu / M^2}{p^2 - m^2 + i\epsilon}$$

Massless Vector Bosons (Photons)

$$-i \frac{g^{\mu\nu}}{p^2}$$

Scalar (Higgs) Bosons

$$i \frac{1}{p^2 - m^2}$$

vertices

B. Appendix B: Numerical Values of Constants

The different factors of vertices are model dependent constants. The numerical values of constants are given by

$$A = e$$

$$B^L = g \left(\frac{\frac{1}{2} - \sin^2 \theta_w}{\sin \theta_w} \right)$$

$$B^R = -g \sin \theta_w$$

$$C = -\frac{g}{\sqrt{2}}$$

$$D = e$$

$$E = g \cos \theta_w$$

$$F_1^L = -g \left(\frac{m_e \cos \alpha}{2M_w \cos \beta} \right)$$

$$F_2^L = g \left(\frac{m_e \cos \alpha}{2M_w \cos \beta} \right)$$

$$F_3^L = 0$$

$$F_1^R = 0$$

$$F_2^R = 0$$

$$F_3^R = ig \left(\frac{m_e \tan \beta}{2M_w} \right)$$

$$G_1 = ig \left[\frac{\sin(\beta - \alpha)}{2 \cos \theta_w} \right]$$

$$G_2 = -ig \left[\frac{\cos(\beta - \alpha)}{2 \cos \theta_w} \right]$$

$$H_1 = gM_w \cos(\beta - \alpha)$$

$$H_2 = gM_w \sin(\beta - \alpha)$$

$$I_1 = \frac{g}{2} \sin(\alpha - \beta)$$

$$I_2 = \frac{g}{2} \cos(\alpha - \beta)$$

$$I_3 = -i\frac{g}{2}$$

$$J_1 = g \left[\frac{M_Z \cos(\beta - \alpha)}{\cos \theta_w} \right]$$

$$J_2 = g \left[\frac{M_Z \sin(\beta - \alpha)}{\cos \theta_w} \right]$$

$$K_L = g \left(\frac{\frac{1}{2} - \sin^2 \theta_w}{\cos \theta_w} \right)$$

$$K_R = -g \left(\frac{\sin^2 \theta_w}{\cos \theta_w} \right)$$

$$L = \frac{g}{\sqrt{2}}$$

$$M = \frac{g}{2 \cos \theta_w}$$

$$N_{Li} = \frac{-g}{2\sqrt{2}} \left(\varepsilon_i \frac{m_e Z_{i3}}{M_w \cos \beta} - Z_{i2} - Z_{i1} \tan \theta_w \right)$$

$$N_{Ri} = \frac{-g}{2\sqrt{2}} \left(\frac{m_e Z_{i3}}{M_w \cos \beta} + 2\varepsilon_i Z_{i1} \tan \theta_w \right)$$

$$N'_{Li} = \frac{-g}{2\sqrt{2}} \left(-\varepsilon_i \frac{m_e Z_{i3}}{M_w \cos \beta} - Z_{i2} - Z_{i1} \tan \theta_w \right)$$

$$N'_{Ri} = \frac{-g}{2\sqrt{2}} \left(\frac{m_e Z_{i3}}{M_w \cos \beta} - 2\varepsilon_i Z_{i1} \tan \theta_w \right)$$

$$O = \frac{g}{\sqrt{2}} (Z_{i1} \tan \theta_w - Z_{i2})$$

$$\begin{aligned}
P_{ij}^L &= \varepsilon_i g \left(Z_{i2} U_{j1} + \frac{Z_{i4} V_{j2}}{\sqrt{2}} \right) \\
P_{ij}^R &= g \left(Z_{i2} U_{j1} + \frac{Z_{i3} U_{j2}}{\sqrt{2}} \right) \\
Q_{ij}^R &= \varepsilon_i g \sin \beta \left[Z_{i3} U_{j1} - \frac{(Z_{i2} + Z_{i1} \tan \theta_w) U_{i2}}{\sqrt{2}} \right]
\end{aligned}$$

$$\begin{aligned}
R_{lij} &= -\frac{g}{2 \sin \beta} \left\{ \frac{m_{\tilde{\chi}_i^0} \delta_{ij} \sin \alpha}{M_w} + (\varepsilon_i + \varepsilon_j) [Q_{ij}'' \sin(\beta - \alpha) - R_{ij}'' \sin \alpha] \right\} \\
R_{2ij} &= -\frac{g}{2 \sin \beta} \left\{ \frac{m_{\tilde{\chi}_i^0} \delta_{ij} \cos \alpha}{M_w} - (\varepsilon_i + \varepsilon_j) [Q_{ij}'' \cos(\beta - \alpha) + R_{ij}'' \cos \alpha] \right\} \\
R_{3ij} &= \frac{ig}{2 \sin \beta} (\varepsilon_i - \varepsilon_j) [Q_{ij}'' \cos 2\beta - R_{ij}'' \cos \beta] \\
R_{1ij}' &= -\frac{g}{2 \sin \beta} [(\varepsilon_i - \varepsilon_j) Q_{ij}'' \sin(\beta - \alpha) - (\varepsilon_i - \varepsilon_j) R_{ij}'' \sin \alpha] \\
R_{2ij}' &= \frac{g}{2 \sin \beta} (\varepsilon_i - \varepsilon_j) [Q_{ij}'' \cos(\beta - \alpha) + R_{ij}'' \cos \alpha] \\
R_{3ij}' &= -\frac{ig}{2 \sin \beta} \left\{ \frac{m_{\tilde{\chi}_i^0} \delta_{ij} \cos \alpha}{M_w} - (\varepsilon_i + \varepsilon_j) [Q_{ij}'' \cos 2\beta + R_{ij}'' \cos \beta] \right\}
\end{aligned}$$

where

$$Q_{ij}'' = \frac{1}{2} [Z_{i3} (g Z_{j2} - g' Z_{j1}) + Z_{j3} (g Z_{i2} - g' Z_{i1})]$$

and

$$R_{ij}'' = \frac{1}{2M_w} [M Z_{i2} Z_{j2} + M' Z_{i1} Z_{j1} - \mu (Z_{i3} Z_{j4} + Z_{i4} Z_{j3})]$$

$$S_{ij} = \frac{g}{4 \cos \theta_w} (1 - \varepsilon_i \varepsilon_j) (Z_{i3} Z_{j3} - Z_{i4} Z_{j4})$$

$$S'_{ij} = \frac{g}{4 \cos \theta_w} (1 + \varepsilon_i \varepsilon_j) (Z_{i3} Z_{j3} - Z_{i4} Z_{j4})$$

$$T_i^L = g \left(\frac{m_e U_{i2}}{\sqrt{2} M_w \cos \beta} \right)$$

$$T_i^R = -g V_{i1}$$

$$U_i = -g U_{i1}$$

$$V=e$$

$$W_{ij}^L = \frac{g}{\cos \theta_w} \left(\delta_{ij} \sin^2 \theta_w - V_{i1} V_{j1} - \frac{1}{2} V_{i2} V_{j2} \right)$$

$$W_{ij}^R = \frac{g}{\cos \theta_w} \left(\delta_{ij} \sin^2 \theta_w - U_{i1} U_{j1} - \frac{1}{2} U_{i2} U_{j2} \right)$$

$$X_{1ij} = -\frac{g}{2 \sin \beta} \left\{ \frac{m_{\tilde{\chi}_i^0} \delta_{ij} \sin \alpha}{M_w} + 2 [Q_{ij} \sin(\beta - \alpha) - R_{ij} \sin \alpha] \right\}$$

$$X_{2ij} = -\frac{g}{2 \sin \beta} \left\{ \frac{m_{\tilde{\chi}_i^0} \delta_{ij} \cos \alpha}{M_w} - 2 [Q_{ij} \cos(\beta - \alpha) + R_{ij} \cos \alpha] \right\}$$

$$X_{3ij} = 0$$

$$X'_{1ij} = 0$$

$$X_{2ij}^{'\prime} = 0$$

$$X_{3ij}^{'\prime} = -\frac{ig}{2\sin\beta} \left\{ \frac{m_{\tilde{\chi}_i^0} \delta_{ij} \cos\beta}{M_w} - 2 [Q_{ij} \cos 2\beta + R_{ij} \cos\beta] \right\}$$

where

$$Q_{ij} = \frac{U_{i2}V_{j1}}{\sqrt{2}}$$

$$R_{ij} = \frac{MU_{i1}V_{j1} + \mu U_{i2}V_{j2}}{2M_w}$$

the values of the constants e , θ_w , g , $g^{'\prime}$, given in natural units ($\hbar = c = 1$) are

$$e = 0.302822$$

$$\theta_w = 0.495293$$

$$g = \frac{e}{\sin\theta_w} = 0.637132$$

$$g^{'\prime} = \frac{e}{\cos\theta_w} = 0.344183$$

The masses of the electron, the Z , and W , bosons are

$$m_e = 0.51099906 \text{ MeV}/c^2$$

$$M_W = 80.33 \text{ GeV}/c^2$$

$$M_Z = 91.187 \text{ GeV}/c^2$$

The width of Z and W bosons are

$$\Gamma_W = 2.07 \text{ GeV}/c^2$$

$$\Gamma_Z = 2.490 \text{ GeV}/c^2$$

The mixing angles α and β and the masses M and M' and μ are model dependent free parameters and they are defined in chapter 3 together with the sign factors η_i and ε_j , masses $m_{\tilde{\chi}_i^\pm}$ and $m_{\tilde{\chi}_i^0}$ and the mixing matrices U , V and Z of charginos and neutralinos.

Three body kinematics

Introduction The energy E and the 3-momentum $\mathbf{p}$ of a particle of mass m form a 4-vector momentum $p \equiv (E, \mathbf{p})$ whose square $p^2 \equiv E^2 - |\mathbf{p}|^2 = m^2$. The scalar dot product of two 4-momentum $p_1 \cdot p_2 = E_1 E_2 - \mathbf{p}_1 \cdot \mathbf{p}_2$ is invariant (frame independent).

In this appendix we describe the three body reaction kinematics and cross-sections in terms of invariant 4-momenta dot products [39], [40].

Dot Products of Momenta

Three body reactions consists of two particles of 4-momenta p_1 and p_2 and masses m_1 and m_2 in the initial state which scatter to three particles of 4-momenta p_3, p_4 , and p_5 and masses m_3, m_4 and m_5 in the final state. When the particles in the initial state are light with respect to their energies (e.g. electrons and positrons); the masses m_1 and m_2 are ignored and we take $m_1=m_2 \simeq 0$.

The Lorentz covariant variables, or the dot products in the case of three body reactions with light particles in the initial state are defined as

$$p_1 \cdot p_1 = m_1^2 \approx 0,$$

$$p_1 \cdot p_2 = \frac{1}{2}(s - m_1^2 - m_2^2) \simeq \frac{s}{2}$$

$$p_1 \cdot p_3 \equiv \zeta$$

$$p_1 \cdot p_4 = \frac{1}{2}(s + m_1^2 - m_2^2 - 2x - 2\zeta) \approx \frac{1}{2}(s - 2x - 2\zeta)$$

$$p_1 \cdot p_5 \equiv x$$

$$p_2 \cdot p_2 = m_2^2 \approx 0$$

$$p_2 \cdot p_3 = \frac{1}{2}(s' + 2y - 2\zeta + m_3^2 - m_4^2)$$

$$\begin{aligned} p_2 \cdot p_4 &= \frac{1}{2}(2\zeta + 2x - 2y - m_1^2 + m_2^2 - m_3^2 + m_4^2 - m_5^2) \\ &\approx \frac{1}{2}(2\zeta + 2x - 2y - m_3^2 + m_4^2 - m_5^2) \end{aligned}$$

$$p_2 \cdot p_5 = \frac{1}{2}(s - s' - 2x + m_5^2)$$

$$p_3 \cdot p_3 = m_3^2$$

$$p_3 \cdot p_4 = \frac{1}{2}(s' - m_3^2 - m_4^2)$$

$$p_3 \cdot p_5 \equiv y$$

$$p_4 \cdot p_4 = m_4^2$$

$$p_4 \cdot p_5 = \frac{1}{2}(s - s' - 2y - m_5^2)$$

$$p_5 \cdot p_5 = m_5^2$$

where the variables s (square of the center of mass energy) and s' are defined from

$$\sqrt{s} = p_1 + p_2$$

$$\sqrt{s'} = p_3 + p_4$$

The function ζ is defined as

$$\zeta = \frac{1}{\Lambda(\sqrt{s}, \sqrt{s'}, m_5^2)} \begin{bmatrix} -m_5^2(s + m_1^2 - m_2^2) \\ +x(s' + m_3^2 - m_4^2)(s - s' + m_5^2) \\ +y(s + m_1^2 - m_2^2)(s - s' - m_5^2) \\ -2xy(s + s') \end{bmatrix}$$

For massless or light particles in the initial state; ζ is approximated as

$$\zeta = \frac{1}{\Lambda(\sqrt{s}, \sqrt{s'}, m_5^2)} \begin{bmatrix} -m_5^2 s(s' + m_3^2 - m_4^2) \\ +x(s' + m_3^2 - m_4^2)(s - s' + m_5^2) \\ +ys(s - s' - m_5^2) \\ -2xy(s + s') \end{bmatrix}$$

where

$$\Lambda(a, b, c) = \sqrt{a^4 + b^4 + c^4 - 2a^2b^2 - 2a^2c^2 - 2b^2c^2}$$

Differential Cross Section

The covariant phase space differential cross-section for three body reactions can be written as

$$d\sigma = \frac{(2\pi)^4 |M|^2}{4\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}} d\Phi_3$$

where M is the matrix element (amplitude) for the reaction or scattering process and $d\Phi_3$ is an element of the three body phase space given by

$$d\Phi_3 = \frac{1}{(2\pi)^9} d\rho_3$$

where $d\rho_3$ is the Lorentz invariant phase space volume element

$$d\rho_3 = \delta^4(p_1 + p_2 - p_3 - p_4 - p_5) \frac{d^3 |\mathbf{p}_3|}{2E_3} \frac{d^3 |\mathbf{p}_4|}{2E_4} \frac{d^3 |\mathbf{p}_5|}{2E_5}$$

the energy and momenta are related by

$$p_i = (E_i, \mathbf{p}_i)$$

$$p_i^2 = E_i^2 - \mathbf{p}_i^2 = m_i^2$$

In center-of-mass frame

$$\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2} = |\mathbf{p}_{lcn}| \sqrt{s}$$

for light particles in the initial state, this expression is approximated as

$$\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2} \approx \sqrt{(p_1 \cdot p_2)^2} = p_1 \cdot p_2 = \frac{1}{2}(s - m_1^2 - m_2^2) \approx \frac{s}{2}$$

Total Cross Section

The Lorentz invariant three body phase space volume element in terms of covariant dot products is given by

$$d\rho_3 = \frac{\pi}{2} \frac{ds' dx dy d\phi}{\Lambda(\sqrt{s}, m_1, m_2) \Lambda(\sqrt{s}, \sqrt{s'}, m_5)}$$

for light particles in the initial state

$$d\rho_3 \approx \frac{\pi}{2} \frac{ds' dx dy d\phi}{\Lambda(\sqrt{s}, 0, 0) \Lambda(\sqrt{s}, \sqrt{s'}, m_5)} = \frac{\pi}{2} \frac{ds' dx dy d\phi}{s \Lambda(\sqrt{s}, \sqrt{s'}, m_5)}$$

where ϕ is the polar angle of $\mathbf{p}_1$ with respect to $\mathbf{p}_5$.

The total cross-section as a function of the center-of-mass energy $\sqrt{s}$ given by

$$\sigma(\sqrt{s}) = \int_{s'_-}^{s'_+} ds' \int_{x_-}^{x_+} dx \int_{y_-}^{y_+} dy \int_{z_-}^{z_+} dz \int_0^{2\pi} d\phi \left[\frac{1}{128\pi^4} \frac{|M|^2}{|\mathbf{p}_{lcn}| \sqrt{s} \Lambda(\sqrt{s}, m_1, m_2) \Lambda(\sqrt{s}, \sqrt{s'}, m_5)} \right]$$

and for light particles in the initial state

$$\sigma(\sqrt{s}) = \int_{s'_-}^{s'_+} ds' \int_{x_-}^{x_+} dx \int_{y_-}^{y_+} dy \int_{z_-}^{z_+} dz \int_0^{2\pi} d\phi \left[\frac{1}{128\pi^4} \frac{|M|^2}{s^2 \Lambda(\sqrt{s}, \sqrt{s'}, m_5)} \right]$$

If the variable angle ϕ is cyclic, the integration over ϕ just contributes a factor of 2π to the phase space. The expression then becomes

$$\sigma(\sqrt{s}) = \int_{s'_-}^{s'_+} ds' \int_{x_-}^{x_+} dx \int_{y_-}^{y_+} dy \left[\frac{1}{64\pi^3} \frac{|M|^2}{s^2 \Lambda(\sqrt{s}, \sqrt{s'}, m_5)} \right]$$

The domain of integration for s' , x , and y are given by

$$s'_- = (m_3 + m_4)^2$$

$$s'_+ = (s - m_5)^2$$

$$\begin{aligned} x_{\pm} &= \frac{1}{4s} \left[(s + m_1^2 + m_2^2) (s - s' + m_5^2) \pm \Lambda(\sqrt{s}, m_1, m_2) \Lambda(\sqrt{s}, \sqrt{s'}, m_5) \right] \\ &\approx \frac{1}{4s} \left[(s + 0 + 0) (s - s' + m_5^2) \pm \Lambda(\sqrt{s}, 0, 0) \Lambda(\sqrt{s}, \sqrt{s'}, m_5) \right] \\ &\approx \frac{1}{4s} \left[(s - s' + m_5^2) \pm \Lambda(\sqrt{s}, \sqrt{s'}, m_5) \right], \\ y_{\pm} &= \frac{1}{4s'} \left[(s' + m_3^2 + m_4^2) (s - s' + m_5^2) \pm \Lambda(\sqrt{s'}, m_3, m_4) \Lambda(\sqrt{s}, \sqrt{s'}, m_5) \right]. \end{aligned}$$

The resulting total cross section is in terms of GeV^{-2} . To convert it into pbarn, we multiply it by the conversion constant

$$(\hbar c)^{-2} = 0.38937966 \times 10^9 GeV^2 \text{ pbarn}.$$

References

1. I. J. Aitchison and A. J. Hey, **Gauge Theories in Particle Physics**, Adam Hilger Ltd., Bristol, 1983.
2. S. M. Bilenky and J. Hosek, Phys. Reports 90 (1982) 73.
3. S. Weinberg, Rev. of Mod. Phys. 52 (1980) 515.
4. A. Salam, Rev. of Mod. Phys. 52 (1980) 525.
5. S. L. Glashow, Rev. of Mod. Phys. 52 (1980) 539.
6. M. A. B. Beg and Sirlin, Phys. Reports 88 (1980) 1.
7. M. Chen et. al., Phys. Reports 159 (1988) 201.
8. M. F. Sohnius, Phys. Reports 128 (1985) 39.
9. H. Baer et. al. Phys. Rev. D 48 (1993) R2978.
10. H. J. W. Mueller-Kirsten and A. Wiedemann, Supersymmetry, World Scientific Ltd., Singapore, 1987;
P. D. B. Collins, Particle Physics and Cosmology, John Wiley & Sons Inc., USA, 1989;
P. Fayet and S. Ferrara, Phys. Reports 32 (1977) 249;
B. Zumino, Phys. Reports 104 (1984) 113;
H. P. Nilles, Phys. Reports 110 (1984) 1.

11. H. E. Haber, et. al., Nucl. Phys. B 161 (1979) 493.
12. A. Mendez, Mod. Phys. Lett. 5 (1990) 2087.
13. P. Fayet, Phys. Reports 105 (1984) 21.
14. H. Georgi, Hadr. Jour. of Phys. 1 (1978) 155.
15. S. Bertolini, Nucl. Phys. B 272 (1986) 77.
16. J. F. Gunion and H. E. Haber, Phys. Rev. D 37 (1988) 2515.
17. A. Bartl et. al., Zeit. f. Phys. C 30 (1986) 441.
18. A. Bartl et. al., Zeit. f. Phys. C 35 (1992) 257.
19. A. Bartl et. al., Nucl. Phys. B 278 (1986) 1.
20. A. Bartl et. al., Phys. Rev. D 40 (1989) 1594.
21. M. M. El-Kheishin, Phys. Rev. D 45 (1992) 4345.
22. H. E. Haber, Phys. Rev. D 54 (1996) 687.
23. E. Ma, Phys. Rev. D 49 (1994) 1663.
24. S. Okubo, Prog. of Theor. Phys. 27 (1962) 1949.
25. R. L. Pease, Inter. Jour. of Theor. Phys. 34 (1995) 183
26. F. Abe et. al., Fermilab Pub. 94/097-E CDF.

- 27. T. Tsukamoto et. al., Phys. Rev. D 51 (1995) 3153.
- 28. J. L. Feng and M. J. Strassler, Phys. Rev. D 51 (1995) 4661.
- 29. J. L. Feng et. al., Phys. Rev. 52 (1995) 1418.
- 30. S. Ambrosanio and B. Mele, Phys. Rev. D 53 (1996) 2541.
- 31. C. Itzykson and J. B. Zuber, Quantum Field Theory, Mc Graw-Hill Inc., USA, 1980.
- 32. M. Veltman, Diagrammatica, Cambridge University Press, 1994.
- 33. H.E. Haber and G. L. Kane, Phys. Rev. D 117 (1985) 75.
- 34. J. F. Gunion and H. E. Haber, Nucl. Phys. B 272 (1986) 1;
J. F. Gunion and H. E. Haber, Nucl. Phys. B 278 (1986) 449;
J. F. Gunion and H. E. Haber, Nucl. Phys. B 307(1988) 445.
- 35. H.E. Haber and D. Wyler, Nucl. Phys. B 323 (1989) 267.
- 36. J. Rosiek, Phys. Rev. D 41 (1990) 3464.
- 37. J. W. Williamson, Am. Jour. of Phys. 33 (1965)987.
- 38. Review of Particle Properties, Phys. Rev. D 54 Part II (1996) 175.
- 39. Alexander Pukhov, CalcHEP 3.2: MSSM, structure functions, event generation, batchs, and generation of matrix elements for other packages, hep-ph/0412191

40. Abdelhak Djouadi, Jean-Loic Kneur, Gilbert Moultaka, SuSpect: a Fortran Code for the Supersymmetric and Higgs Particle Spectrum in the MSSM, hep-ph/0211331